

Small deformations of polygons

Jean-Marc Schlenker*

October 2004; revised, December 2004 (v2)

Abstract

We describe the first-order variations of the angles of Euclidean, spherical or hyperbolic polygons under infinitesimal deformations such that the lengths of the edges do not change. Using this description, we introduce a vector-valued quadratic invariant b on the space of those isometric deformations which, for convex polygons, has a remarkable positivity property.

We give two geometric applications. The first is an isoperimetric statement for hyperbolic polygons: among the convex hyperbolic polygons with given edge lengths, there is a unique polygon with vertices on a circle, a horocycle, or on one connected component of the space of points at constant distance from a geodesic, and it has maximal area. The second application is a new proof of the infinitesimal rigidity of convex polyhedra in the Euclidean space, and a new rigidity result for polyhedral surfaces in the Minkowski space.

Finally we indicate how the invariant b can be used to define natural metrics on the space of convex spherical (or hyperbolic) polygons with fixed edge lengths. Those metrics are related to known (and interesting) metrics on the space of convex polygons with given angles in the plane.

Résumé

On décrit les déformations infinitésimales des angles d'un polygone euclidien, sphérique ou hyperbolique sous les déformations infinitésimales qui préservent les longueurs des arêtes. On en déduit la définition d'un invariant quadratique à valeurs vectorielles b sur l'espace de ces déformations isométriques qui, pour les polygones convexes, a une propriété remarquable de positivité.

On donne deux applications géométriques. La première est un énoncé isopérimétrique pour les polygones hyperboliques: parmi les polygones hyperboliques convexes dont les longueurs des arêtes sont données, il existe un unique élément dont les sommets sont sur un cercle, un horocycle, ou dans une composante connexe de l'ensemble des points à distance constante d'une droite, et son aire est maximale. La seconde application est une nouvelle preuve de la rigidité infinitésimale des polyèdres euclidiens, et un nouveau résultat de rigidité pour les surfaces polyédrales dans l'espace de Minkowski.

Finalement on indique comment l'invariant b peut être utilisé pour définir des métriques naturelles sur l'espace des polygones convexes sphériques (ou hyperboliques) dont les longueurs des cotés sont fixés. Ces métriques sont reliées à des métriques connues, et intéressantes, sur les espaces de polygones euclidiens convexes dont les angles sont fixés.

Contents

1	Introduction	2
2	Deformations of polygons	7
3	A positivity result	9
4	Isoperimetric problems for spherical or hyperbolic polygons	15
5	Infinitesimal rigidity of Euclidean polyhedra	21

*Laboratoire Emile Picard, UMR CNRS 5580, UFR MIG, Université Paul Sabatier, 118 route de Narbonne, 31062 Toulouse Cedex 4, France. schlenker@picard.ups-tlse.fr; <http://picard.ups-tlse.fr/~schlenker>.

1 Introduction

Moduli spaces of polygons. In this paper, a polygon with n vertices in the Euclidean plane E^2 (resp. the sphere S^2 , the hyperbolic plane H^2 or the de Sitter plane S_1^2) is a finite sequence of points $v_1, v_2, \dots, v_n$ in E^2 (resp. S^2 , H^2 , S_1^2), with $v_0 := v_n$, and with $v_i \neq v_{i+1}$ for all $i \in \{1, \dots, n\}$. We call $\mathcal{P}_{n,E}$ (resp. $\mathcal{P}_{n,S}$, $\mathcal{P}_{n,H}$, $\mathcal{P}_{n,dS}$) the moduli space of polygons in each of the spaces, i.e. the space of polygons quotiented by the group of orientation-preserving isometries of the space. In the sphere and the de Sitter plane, we only consider polygons such that two consecutive vertices are never antipodal (in the sphere, they are at distance less than π) although those cases could be included at the cost of a little more care.

The edge lengths of a polygon are the numbers $l_1, \dots, l_n$, where l_i is the distance between v_i and v_{i+1} . Given $l = (l_1, \dots, l_n)$, $\mathcal{P}_E(l)$ is the subset of $\mathcal{P}_{n,E}$ of Euclidean polygons with edge lengths equal to $l_1, \dots, l_n$. Similar notations will be used for spherical, hyperbolic or de Sitter polygons.

First-order deformations of polygons. Let $p = (v_1, \dots, v_n)$ be a Euclidean polygon, and let $\alpha_1, \dots, \alpha_n$ be its angles. An isometric first-order deformation of p is a set of vectors $\dot{v}_1, \dots, \dot{v}_n$, with $\dot{v}_i \in T_{v_i}E^2$, such that, if one deforms p infinitesimally by moving each v_i along $\dot{v}_i$, the lengths of the edges do not change (at first order). This is equivalent to the fact that, for each $i \in \{0, 1, \dots, n-1\}$, $\langle v_{i+1} - v_i, \dot{v}_{i+1} - \dot{v}_i \rangle = 0$. An infinitesimal first-order deformation of p is *trivial* if there exists a Killing field κ on $\mathbb{R}^2$ such that, for each i , $\dot{v}_i = \kappa(v_i)$. We mostly consider the isometric first-order deformations of p up to the trivial deformations; for a “generic” polygon p , they are canonically associated to the elements of $T_{[p]}\mathcal{P}_E(l)$, where $l = (l_1, \dots, l_n)$ is the set of lengths of p and $[p]$ is the image of p under the projection from the space of polygons in $\mathbb{R}^2$ of given edge lengths to the quotient under the action of $\text{Isom}(\mathbb{R}^2)$.

Clearly, the isometric first-order deformations of p are determined by the first-order variations $\dot{\alpha}_i$ of its angles. However not all possible variations are possible. We first give some constraints on the possible first-order variations of the angles of polygons in the spaces we consider.

Proposition 1.1. *Let $p = (v_1, \dots, v_n)$ be a Euclidean polygon. Let $\alpha_1, \dots, \alpha_n$ be its angles, and let $\dot{\alpha}_1, \dots, \dot{\alpha}_n \in \mathbb{R}$. Suppose that there exists a first-order isometric deformation of p such that the induced first-order variation of the angles of p is $(\dot{\alpha}_1, \dots, \dot{\alpha}_n)$. Then:*

- $\sum_{i=1}^n \dot{\alpha}_i = 0$.
- $\sum_{i=1}^n \dot{\alpha}_i v_i = 0$.

Conversely, if those two conditions are satisfied and the vertices of p are not all on a line, then there exists an isometric first-order deformation of p such that the induced first-order variation of the α_i are the $\dot{\alpha}_i$, and it is unique up to the addition of a trivial deformation.

An analogous but even simpler statement holds for spherical polygons.

Theorem A_S. *Let $p = (v_1, \dots, v_n)$ be a spherical polygon. Let $\alpha_1, \dots, \alpha_n$ be its angles, and let $\dot{\alpha}_1, \dots, \dot{\alpha}_n \in \mathbb{R}$ be a first-order variation of its angles induced by an isometric first-order deformation of p . Then:*

$$\sum_{i=1}^n \dot{\alpha}_i v_i = 0 ,$$

where the v_i are considered as points in $S^2 \subset \mathbb{R}^3$.

Conversely, if this equation is satisfied by an n -uple $(\dot{\alpha}_1, \dots, \dot{\alpha}_n)$ and moreover the v_i are not all on a great circle, then there exists an isometric first-order deformation of p such that the $\dot{\alpha}_i$ are the associated first-order variations of the α_i .

Each first-order isometric deformation of p is uniquely determined, up to the addition of a trivial deformation, by the corresponding variation of its angles. Moreover, the space of isometric first-order deformations of p (up to the trivial ones) has dimension at least $n - 3$, since a deformation of p is determined by $2n$ parameters (the positions of the vertices), with n constraints (the lengths of the edges) and a group of isometries of dimensions 3. Therefore, the space of first-order variations of the α_i , induced by isometric first-order deformations of p , always has dimension at least $n - 3$. It follows that, unless the v_i are on a great circle, the space of isometric first-order deformations of p has dimension $n - 3$. We will see at the beginning of section 2 that this implies the fact, proved by Kapovich and Millson [KM99], that in this case the space of polygons with the same edge lengths as p is locally a smooth manifold.

The same description holds in the hyperbolic plane.

Theorem A_H . *Let $p = (v_1, \dots, v_n)$ be a hyperbolic polygon. Let $\alpha_1, \dots, \alpha_n$ be its angles, and let $\dot{\alpha}_1, \dots, \dot{\alpha}_n \in \mathbb{R}$ be a first-order variation of its angles induced by an isometric first-order deformation of p . Then:*

$$\sum_{i=1}^n \dot{\alpha}_i v_i = 0 ,$$

where the v_i are considered as points in $H^2 \subset \mathbb{R}_1^3$.

Conversely, if this equation is satisfied by an n -uple $(\dot{\alpha}_1, \dots, \dot{\alpha}_n)$ and moreover the v_i are not all on a hyperbolic geodesic, then there exists an isometric first-order deformation of p such that the $\dot{\alpha}_i$ are the associated first-order variations of the α_i .

Here $\mathbb{R}_1^3$ is the Minkowski 3-dimensional space, i.e. it is $\mathbb{R}^3$ with the bilinear form:

$$\langle (x, y, z), (x', y', z') \rangle = xx' + yy' - zz' .$$

H^2 has an isometric embedding in $\mathbb{R}_1^3$ as a quadric:

$$H^2 = \{(x, y, z) \in \mathbb{R}_1^3 \mid \langle (x, y, z), (x, y, z) \rangle = -1 \text{ and } z > 0\} .$$

A quadratic form on first-order deformations. This description of the first-order deformations of the angles of polygons opens the door to the definition of a quadratic invariant of first-order isometric deformations.

Definition 1.2. *Let $p = (v_1, \dots, v_n)$ be an Euclidean polygon, with edge lengths $l_1, \dots, l_n$. Let $l := (l_1, \dots, l_n)$, and let $U \in T_{[p]} \mathcal{P}_E(l)$, where $[p]$ is the projection of p in $\mathcal{P}_E(l)$. Let U' be a first-order deformation of p projecting, under the quotient by the trivial deformations, to U . We call:*

$$b(U) := \sum_{i=1}^n d\alpha_i(U') dv_i(U') ,$$

where the $d\alpha_i(U')$ and the $dv_i(U')$ are the first-order variations of the α_i and of the v_i , respectively, under U' . Then $b(U)$ does not depend on the choice of U' .

As a first-order deformation of p defined up to the trivial deformations, U does not define uniquely the first-order variations of the v_i . However, any choice U' of a representative $(\dot{v}_1, \dots, \dot{v}_n)$ will do. Indeed, let U'' be another first-order deformation of p corresponding to U ; then $U' - U''$ is a trivial deformation, so the corresponding variation of the v_i is given by $\kappa(v_i)$, where κ is a Killing field. But any Killing field κ in E^2 is of the form:

$$\kappa(x) = e^{i\theta} x + \tau ,$$

for some $\theta \in \mathbb{R}$ and some vector $\tau \in E^2$ (this expression uses the usual identification of E^2 with $\mathbb{C}$). It follows that:

$$\sum_{i=1}^n d\alpha_i(U') dv_i(U') - \sum_{i=1}^n d\alpha_i(U'') dv_i(U'') = \sum_{i=1}^n d\alpha_i(U') \kappa(v_i) =$$

$$= \sum_{i=1}^n d\alpha_i(U')(e^{i\theta}v_i + \tau) = e^{i\theta} \left(\sum_{i=1}^n d\alpha_i(U')v_i \right) + \left(\sum_{i=1}^n d\alpha_i(U') \right) \tau ,$$

and both terms vanish by Proposition 1.1.

A very similar definition can be used in the sphere or the hyperbolic plane.

Definition 1.3. Let $p = (v_1, \dots, v_n)$ be a polygon in the sphere (resp. the hyperbolic plane) with edge lengths $l_1, \dots, l_n$. Let $l := (l_1, \dots, l_n)$, and let $U \in T_{[p]}\mathcal{P}_S(l)$ (resp. $T_{[p]}\mathcal{P}_H(l)$) where $[p]$ is the projection of p in $\mathcal{P}_S(l)$ (resp. $\mathcal{P}_H(l)$). Let U' be a first-order deformation of p projecting, under the quotient by the trivial deformations, to U . We call:

$$b(U) := \sum_{i=1}^n d\alpha_i(U')dv_i(U') ,$$

where the $d\alpha_i(U')$ and the $dv_i(U')$ are the first-order variations of the α_i and of the v_i , considered as points in $\mathbb{R}^3$ (resp. $\mathbb{R}_1^3$), respectively, under U' . Then $b(U)$ does not depend on the choice of U' .

An argument very similar to the one given above shows that $b(U)$ is indeed independent of the precise deformation U' of p which is chosen. In the sphere, any Killing field is of the form:

$$\kappa(x) = Y \times x ,$$

for some vector $Y \in \mathbb{R}^3$. Therefore, if U'' is another first-order deformation of p projecting to U , then $U' - U''$ corresponds to a Killing field κ . Then:

$$\begin{aligned} \sum_{i=1}^n d\alpha_i(U')dv_i(U') - \sum_{i=1}^n d\alpha_i(U'')dv_i(U'') &= \sum_{i=1}^n d\alpha_i(U')\kappa(v_i) = \\ &= \sum_{i=1}^n d\alpha_i(U')Y \times v_i = Y \times \left(\sum_{i=1}^n d\alpha_i(U')v_i \right) = 0 . \end{aligned}$$

The same argument can be used in the hyperbolic context, using the fact that the Killing fields of $\mathbb{R}_1^3$ which vanish at the origin (in other terms the elements of the Lie algebra $so(2,1)$) are of the form:

$$\kappa(x) = Y \boxtimes x ,$$

where $\boxtimes$ is the Minkowski analog of the vector product:

$$(Y_1, Y_2, Y_3) \boxtimes (x_1, x_2, x_3) = (Y_2x_3 - Y_3x_2, Y_3x_1 - Y_1x_3, -Y_1x_2 + Y_2x_1) .$$

In both cases, the fact that b depends only on the equivalence class of U' (under the action of trivial deformations) has an “abstract” interpretation. As defined above, for each polygon p in S^2 , b defines a quadratic form on the space of isometric first-order deformations of p , with values in $\mathbb{R}^3$. Let $l = (l_1, \dots, l_n)$ be the edge lengths of p . Since b behaves “well” under the action of $SO(3)$ on S^2 , b actually defines a quadratic form on $T_{[p]}\mathcal{P}_S(l)$, where $[p]$ is the projection of p in $\mathcal{P}_S(l)$, with values in the vector bundle over $\mathcal{P}_S(l)$ which is defined, from the trivial $\mathbb{R}^3$ -bundle over the space of polygons in S^2 with edge lengths given by l , by taking the natural action of $SO(3)$ on both this space of polygons and $\mathbb{R}^3$.

A positivity property. The quadratic form b defined above has a striking geometric property when p is a convex spherical or hyperbolic polygon. By a *convex polygon*, we mean a polygon which is the boundary of a convex domain in S^2 (resp. H^2), which we call the *interior* of p , and denote by $\text{int}(p)$. We state this property first in the spherical setting. Recall that, given a convex spherical polygon p , the *dual polygon* p^* is a convex polygon whose interior is the set of points in S^2 which have positive scalar product with the vertices of p (see e.g. [Cox93, Cox57]). Each of its vertices is at distance $\pi/2$ of an edge of p , and conversely.

Theorem B_S. *Let p be a convex polygon in S^2 , and let U be a non-trivial infinitesimal first-order deformation of p . Then $b(U) \in (\mathbb{R}_+ \setminus \{0\})\text{int}(p^*)$, i.e. $b(U)$ is contained in the positive cone over the interior of p^* .*

There is a geometric interpretation to this property. The space of polygons with given edge lengths $l = (l_1, \dots, l_n)$, $\mathcal{P}_S(l)$, has a natural map ϕ to $\mathbb{R}^n$, where $\phi(p)$ is the family of the angles of p . By Theorem A_S, the image is a locally a submanifold except when p has all its vertices on a great circle. We will see in section 3 that b is strongly related to the second fundamental form of this submanifold, so that Theorem B_S translates as a convexity property: if p is a convex polygon, the image of ϕ , in the neighborhood of $\phi(p)$, has a second fundamental form which is positive definite in some directions. Theorem B_S is related to a result of Volkov [Vol56] on (non infinitesimal) isometric deformations of spherical polygons.

The same result holds in the hyperbolic or the de Sitter setting. Now the dual of a hyperbolic polygon is a convex de Sitter polygon, and conversely; both can be defined as in the sphere, using the Minkowski metric on $\mathbb{R}_1^3$ (see e.g. [Cox43, Cox57], where the “polarities” corresponding to the duality used here has a very central role).

Theorem B_H. *Let p be a convex polygon in H^2 and let U be a non-trivial infinitesimal first-order deformation of P . Then $b(U) \in (\mathbb{R}_+ \setminus \{0\})\text{int}(p^*)$, i.e. $b(U)$ is in the positive cone over the interior of p^* .*

An isoperimetric statement for hyperbolic polygons. The first application of Theorems A_H and B_H that we consider is to an isoperimetric problem for hyperbolic polygons. The next theorem is hyperbolic analog of a statement which was proved by Steiner [Ste42] in the spherical setting, but was known earlier in the Euclidean case (see [Sie02] for recent progress on more elaborate statements even in the Euclidean case).

Given $l = (l_1, \dots, l_n)$, we call $\mathcal{P}_S^c(l)$ the space of convex polygons in S^2 with edge lengths equal to the l_i , considered up to global isometries in S^2 , and $\mathcal{P}_H^c(l)$ the space of convex hyperbolic polygons with edge lengths equal to the l_i , again considered up to global isometries.

Theorem 1.4. *Let $l = (l_1, \dots, l_n)$ be such that $\mathcal{P}_H^c(l)$ is not empty; then there exists a unique element $p \in \mathcal{P}_H^c(l)$ which has its vertices either on a circle, on a horocycle, or on a connected set of points at fixed distance from a geodesic. Moreover, p has maximal area.*

There is a similar statement, which is slightly simpler, in the Euclidean plane and in the sphere; in both cases, there is a unique polygon of maximal area (among the polygons with fixed edge lengths) and it has its vertices on a circle. The spherical result was apparently discovered by Steiner [Ste42], but he writes that the Euclidean statement was previously known. The proof given here for hyperbolic polygons also works in the spherical setting.

The first point of the proof is that, as a direct consequence of Theorem A_H, a hyperbolic polygon is a critical point of the area, restricted to polygons with the same edge lengths, if and only if its vertices are either on a circle, a horocycle, or a connected component of the space of points at fixed distance from a geodesic.

Theorem B_S (resp. B_H) also has an interesting meaning in this context: it implies that the area, as a function defined on the space of convex polygon with given edge lengths, is “often” strictly concave for a natural metric.

The second point of the proof is that, on the boundary of the space of convex polygons with given edge lengths, the interior normal derivative of the area is positive. It follows that there exists at least one local maximum of the area in the interior. But a direct argument shows that, given the edge lengths, there is at most one polygon with its vertices on a circle, a horocycle, or on one connected component of the set of points at fixed distance from a geodesic.

The infinitesimal rigidity of convex polyhedra. The positivity property of b in the spherical setting leads to a simple proof of the infinitesimal rigidity of convex polyhedra in the Euclidean space. Although the first proof was given by Dehn [Deh16], it follows from the ideas of Legendre [LegII] and Cauchy [Cau13]. Other proofs have been given, in particular by Kann [Kan90] and Filliman [Fil92].

Theorem C. (Legendre, Cauchy, Dehn) *Let P be a convex polyhedron in $\mathbb{R}^3$. Any first-order deformation of P which does not change its combinatorics or the metrics on its faces is trivial, i.e. induced by a global Killing field.*

The proof given here is in section 5. It bears some relations with a proof of the global rigidity of convex Euclidean polyhedra discovered by Pogorelov [Pog56].

Fuchsian polyhedral surfaces in the Minkowski space. The techniques described here have a natural application to another rigidity problem, concerning polyhedral surfaces in the Minkowski 3-dimensional space. We consider space-like polyhedra; rather than closed polyhedra, which can be defined as images of convex polyhedral maps from the sphere, we consider *equivariant polyhedra*, which are the images of a polyhedral map from the universal cover of a surface of genus at least 2 which is *equivariant* (more precise definitions can be found in section 6). We will prove that, among those surfaces, those which are “Fuchsian” – the associated representation from the fundamental group of the surface to the isometry group of $\mathbb{R}_1^3$ has its image in $SO(2, 1)$ – are infinitesimally rigid, i.e. any first-order deformation of those surfaces which does not change the induced metric is trivial, this is Theorem 6.2.

There is an analogous result for smooth, equivariant, Fuchsian, convex surfaces [LS00], which was proved by related methods. Actually, finding a polyhedral version of the results of [LS00] was the main motivation for the present work, although the by-products turned out to be rather more interesting. This question was also studied by I. Iskhakov [Isk00], who provided some partial results. One feature of the proof of the rigidity of convex polyhedra given here is the existence of a distinguished point, which appears quite artificial in the Euclidian context; for surfaces in the Minkowski space, however, it is quite natural and even necessary, since the point which is fixed by the representation of the fundamental group already plays a special role.

The infinitesimal rigidity of Fuchsian equivariant surfaces in the Minkowski space is equivalent, thanks to the “Pogorelov map” used e.g. in [LS00], to similar rigidity statements in the de Sitter or anti-de Sitter space, so that polyhedral rigidity results in those spaces could be obtained as a consequence of the rigidity theorem for polyhedral surfaces in $\mathbb{R}_1^3$ proved here. However, another proof of the rigidity of Fuchsian equivariant surfaces in the de Sitter or the anti-de Sitter space has recently been developed by F. Fillastre (in preparation), and it applies in particular in the polyhedral setting. So it should also be possible to prove the result stated here – on polyhedral Fuchsian surfaces in $\mathbb{R}_1^3$ – from the statements obtained by Fillastre in the de Sitter or the anti-de Sitter space.

To prove Theorem 6.2, we will follow the proof of Theorem C, but the polygons that will be considered will be in the de Sitter plane rather than in the sphere. Recall that, in addition to the hyperbolic plane, $\mathbb{R}_1^3$ contains another quadric, the de Sitter plane S_1^2 , which is a complete, constant curvature 2-dimensional Lorentz manifold (see e.g. [Cox43, Cox57, O’N83]):

$$S_1^2 = \{(x, y, z) \in \mathbb{R}_1^3 \mid \langle (x, y, z), (x, y, z) \rangle = 1\} .$$

Polygons in the de Sitter plane can be defined as in the sphere; the edges can be of different types, either space-like, light-like or time-like. The notion of angle is more subtle than in the sphere, see e.g. [Sch98, Sch01], and it is quite natural to consider the angles of a polygon as complex numbers, with real part either $\pm\pi/2$ or π . A key point is that, with those definitions and the corresponding definitions for the lengths of the edges, the main triangle formulas in the sphere remain valid in the de Sitter plane, a simple fact which we prove in section 6 for completeness.

The first-order deformations of polygons in the de Sitter plane, in terms of the first-order variation of the angles, can be described as in the sphere or the hyperbolic plane.

Theorem A_{ds}. *Let $p = (v_1, \dots, v_n)$ be a polygon in the de Sitter plane. Let $\alpha_1, \dots, \alpha_n$ be its angles, and let $\dot{\alpha}_1, \dots, \dot{\alpha}_n \in \mathbb{R}$ be a first-order variation of its angles induced by an isometric first-order deformation of p . Then:*

$$\sum_{i=1}^n \dot{\alpha}_i v_i = 0 ,$$

where the v_i are considered as points in $\mathbb{R}_1^3$.

Conversely, if this equation is satisfied by an n -uple $(\dot{\alpha}_1, \dots, \dot{\alpha}_n)$ and moreover the v_i are not all on geodesic, then there exists an isometric first-order deformation of p such that the $\dot{\alpha}_i$ are the associated first-order variations of the α_i .

In this context, the quadratic form b can be defined as in the sphere. It has the same positivity property for convex, space-like polygons (a notion which is defined here with some care, see section 6). The notion of polygon dual to a space-like, convex polygon is defined in section 6 as in the sphere, it is a polygon in the hyperbolic plane.

Theorem B_{ds}. *Let p be a convex, space-like polygon in S_1^2 and let U be an infinitesimal first-order deformation of P . Then $b(U) \in (\mathbb{R}_+ \setminus \{0\})\text{int}(p^*)$, i.e. $b(U)$ is contained in the positive cone over the interior of the dual polygon p^* .*

The argument used in the proof of Theorem C, translated to the Minkowski setting, yields a rigidity statement for Fuchsian polyhedral surfaces in $\mathbb{R}_1^3$, Theorem 6.2.

Metrics on moduli spaces. One consequence of Theorem B_S is that it shows that the invariant b can be used to define some natural metrics on the moduli space of convex polygons in S^2 (and similarly in the hyperbolic plane). This is described in section 4, along with some relations to a “natural” metric on Euclidean polygons with fixed angles, which has some interesting properties.

Notations. In all the paper, we set: $\mathbb{R}_+^* := \mathbb{R}_+ \setminus \{0\}$, and $\mathbb{R}_-^* := \mathbb{R}_- \setminus \{0\}$

2 Deformations of polygons

The moduli space of polygons. The geometry of the moduli space of polygons has been studied rather extensively, in particular by Kapovich and Millson (see e.g. [KM95, KM99]). We recall here only some very elementary properties which should clarify parts of the proofs below. We consider here polygons in the plane, however all the comments in this paragraph hold also for spherical, hyperbolic or de Sitter polygons, with some obvious adaptations.

Let $n \geq 3$, let $P_{n,E}$ be the space of polygons with n vertices in $\mathbb{R}^2$. Recall that a polygon is a sequence of vertices $v_1, \dots, v_n = v_0$ such that, for each i , $v_i \neq v_{i+1}$. The isometry group $\text{Isom}(\mathbb{R}^2)$ of $\mathbb{R}^2$ acts without fixed points on $P_{n,E}$, so we consider the quotient $\mathcal{P}_{n,E} = P_{n,E}/\text{Isom}(\mathbb{R}^2)$. $\mathcal{P}_{n,E}$ is a smooth manifold of dimension $2n - 3$, which can also be considered as a smooth algebraic variety.

There is a family of n functions naturally defined on $\mathcal{P}_{n,E}$; if $p \in P_{n,E}$ is a polygon, with $p = (v_1, \dots, v_n)$, then:

$$\lambda_i(p) := d(v_i, v_{i+1}) .$$

Those n functions are clearly invariant under the action of $\text{Isom}(\mathbb{R}^2)$, so they define natural functions on $\mathcal{P}_{n,E}$, which we still call λ_i . Note that it would be algebraically more natural to consider the *squares* of the distances, but we stick to the more natural definition from an elementary geometry viewpoint (for spherical polygons one could consider the cosine of the distance, and for hyperbolic polygons the cosh).

Now let $(l_1, \dots, l_n)$ be an n -uple of positive numbers, recall that $\mathcal{P}_E(l)$ is the (moduli) space of polygons with edge lengths l_i . So:

$$\mathcal{P}_E(l) := \{p \in \mathcal{P}_{n,E} \mid \forall i \in \{1, \dots, n\}, \lambda_i(p) = l_i\} ,$$

so that $\mathcal{P}_E(l)$ is an algebraic subvariety of $\mathcal{P}_{n,E}$. Let $p \in \mathcal{P}_E(l)$, consider the space of its first-order infinitesimal deformations:

$$\mathcal{T}_p := \ker(d\lambda_1) \cap \dots \cap \ker(d\lambda_n) \subset T_p \mathcal{P}_{n,E} .$$

Clearly $\mathcal{T}_p$ has dimension at least $n - 3$, since it is defined by n equations. If those n equations are linearly independent, then, by the inverse function theorem, in the neighborhood of p , $\mathcal{P}_E(l)$ is a smooth submanifold of $\mathcal{P}_{n,E}$ (and also a smooth algebraic subvariety) of dimension $n - 3$.

Euclidean polygons. We first indicate the proof of Proposition 1.1. The simplest proof is perhaps obtained by taking a limit, in Theorem A_S below, when the lengths of the edges go to 0. It is however possible to give a simpler direct proof (this was pointed out by Sergiu Moroianu).

Let $p = (v_1, \dots, v_n)$ be a Euclidean polygon, with edge lengths $l_1, \dots, l_n$, so that, for each $i \in \{1, \dots, n\}$, $l_i = \|v_{i+1} - v_i\| \neq 0$ (with $v_n = v_0$). For each i , let θ_i be the (oriented) angle between the oriented x -axis and $v_{i+1} - v_i$. The possible values of the angles $\theta_1, \dots, \theta_n$, in the neighborhood of p , are defined by condition that $\sum_{i=1}^n l_i e^{i\theta_i} = 0$, because $v_{n+1} = v_1$.

In an isometric first-order deformation of p , it follows that:

$$\sum_{i=1}^n \dot{\theta}_i l_i e^{i\theta_i} = 0 .$$

Suppose (adding a trivial deformation if necessary) that, in the deformation of p we consider, the direction of $v_1 - v_0$ does not vary. Then, for each i , $\dot{\theta}_i = \sum_{j=1}^i \dot{\alpha}_j$. Therefore:

$$0 = \sum_{i=1}^n l_i e^{i\theta_i} \sum_{j=1}^i \dot{\alpha}_j = \sum_{j=1}^n \dot{\alpha}_j \sum_{i=j}^n l_i e^{i\theta_i} .$$

We can also suppose – still without loss of generality – that $v_0 = 0$; then, for each j :

$$v_j = \sum_{i=0}^{j-1} l_i e^{i\theta_i} = - \sum_{i=j}^n l_i e^{i\theta_i} ,$$

and it follows that:

$$\sum_{j=1}^n \dot{\alpha}_j v_j = 0 .$$

In addition, it is well known that, for any Euclidean polygon:

$$\sum_{i=1}^n \alpha_i = 2\pi k ,$$

where k is the winding number of the polygon. It follows that, in any first-order deformation, $\sum_{i=1}^n \dot{\alpha}_i = 0$.

The two conditions in the statement of Proposition 1.1 are linearly independent if and only if the v_i are not all on a line. When the v_i are not collinear, the vector space of n -uples $(\dot{\alpha}_1, \dots, \dot{\alpha}_n)$ satisfying them has dimension $n - 3$. Therefore, the arguments in the previous paragraph show that this space has dimension exactly $n - 3$, and the possible first-order variations of the α_i are exactly the solutions of the two equations in the statement of the proposition.

Spherical polygons. We now prove Theorem A_S . A direct proof is possible¹, but the proof given here is geometric and much simpler.

Let $p = (v_1, \dots, v_n)$ be a spherical polygon, and let U be a first-order isometric deformation of p . Consider the polyhedral cone C over p , i.e. the union of the half-lines with endpoint at 0 which intersect $p \subset S^2 \subset \mathbb{R}^3$. C has n faces $f_1, \dots, f_n$, with f_i equal to the cone over the edge e_i of p . Since α_i is the angle between the edges e_i and e_{i+1} of p , it is also equal to the dihedral angle between f_i and f_{i+1} .

The first-order deformation U of P determines a first-order deformation V of C . For each $i \in \{1, \dots, n\}$, V acts on each face of C as a Killing field, i.e. there exist vectors u_i and y_i such that V acts on f_i as the vector field κ_i defined by:

$$\forall x \in \mathbb{R}^3, \kappa_i(x) = u_i + y_i \times x .$$

Then $\kappa_{i+1} - \kappa_i = 0$ on the intersection of f_i and f_{i+1} , and the difference between κ_i and κ_{i+1} is equal to an infinitesimal rotation of angle $d\alpha_i(U)$ and axis $f_i \cap f_{i+1}$. It follows that:

$$y_{i+1} = y_i + d\alpha_i(U) v_i ,$$

and the result follows since the sum of the differences $y_{i+1} - y_i$, for i going from 1 to n , vanishes.

¹One such proof is available in the first version of this paper, see <http://front.math.ucdavis.edu/math.DG/0410058> and then choose “v1”.

Hyperbolic polygons. The same argument can be used to prove Theorem A_H . C is now a polyhedral cone in the Minkowski space $\mathbb{R}_1^3$, and its faces are time-like. The restriction of the infinitesimal deformation V to each face f_i is a Killing vector, which is now of the form: $\kappa_i(x) = u_i + y_i \boxtimes x$. The proof then proceeds as for the sphere.

3 A positivity result

In this section we prove Theorem B_S , as well as its hyperbolic analog Theorem B_H , after some preliminary computations concerning the first-order deformations of quadrilaterals.

Deformations of quadrilaterals. We consider here a quadrilateral $q = (v_1, v_2, v_3, v_4)$ for which no 3 of the vertices are collinear, and set $v_0 := v_4$. We use the notations apparent in Figure 3, in particular t is the distance between v_1 and v_3 . We also call $d_{i,j}$ the distance between v_i and v_j .




Figure 1: Deformation of a non-degenerate quadrilateral.

Consider a first-order isometric deformation of q , suppose for instance that v_0 and v_1 are fixed (this can be achieved by adding a trivial deformation). Then v_2 moves in the direction of the circle of center v_1 containing it. Since v_1, v_2 and v_3 are not collinear, the circle of center v_3 containing v_2 is not tangent to the circle of center v_1 containing v_2 , so that any first-order displacement of v_2 induces a non-zero first-order variation of the distance from v_2 to v_3 .

The same argument shows that any first-order displacement of v_3 , preserving its distance to v_0 , induces a non-zero first-order variation of its distance to v_2 . It follows that there is a 1-dimensional vector space of isometric first-order deformations of q (up to the trivial deformations), they are parametrized for instance by t , the distance from v_1 to v_3 . So we consider the first-order deformation U of q such that $dt(U) = 1$.

To simplify notations, we call d_{ij} the distance between v_i and v_j , for $0 \leq i, j \leq 4$. Then a well-known spherical geometry formula states that:

$$\cos(t) = \cos(d_{01}) \cos(d_{03}) + \cos(\alpha_0) \sin(d_{01}) \sin(d_{03}) ,$$

so that:

$$d\alpha_0(U) = \frac{\sin(t)}{\sin(d_{01}) \sin(d_{03}) \sin(\alpha_0)} .$$

The same computation (or a symmetry argument) shows that:

$$d\alpha_2(U) = \frac{\sin(t)}{\sin(d_{12}) \sin(d_{23}) \sin(\alpha_2)} .$$

Moreover, equation (1), applied to the triangle (v_0, v_1, v_3) , yields:

$$\cos(\beta_1) = \frac{\cos(d_{03}) - \cos(d_{01}) \cos(t)}{\sin(d_{01}) \sin(t)} ,$$

so that:

$$\begin{aligned} -\sin(\beta_1)d\beta_1(U) &= \frac{\cos(d_{01})\sin^2(t) - (\cos(d_{03}) - \cos(d_{01})\cos(t))\cos(t)}{\sin(d_{01})\sin^2(t)} \\ &= \frac{\cos(d_{01}) - \cos(d_{03})\cos(t)}{\sin(d_{01})\sin^2(t)}. \end{aligned}$$

But, by the sine formula for spherical triangles:

$$\frac{\sin(\beta_1)}{\sin(d_{03})} = \frac{\sin(\alpha_0)}{\sin(t)},$$

so that:

$$d\beta_1(U) = \frac{\cos(d_{03})\cos(t) - \cos(d_{01})}{\sin(d_{01})\sin(d_{03})\sin(t)\sin(\alpha_0)}.$$

The same computation (or a symmetry argument) shows that:

$$d\gamma_1(U) = \frac{\cos(d_{23})\cos(t) - \cos(d_{12})}{\sin(d_{12})\sin(d_{23})\sin(t)\sin(\alpha_2)},$$

and, taking the sum, we obtain that:

$$d\alpha_1(U) = \frac{\cos(d_{03})\cos(t) - \cos(d_{01})}{\sin(d_{01})\sin(d_{03})\sin(t)\sin(\alpha_0)} + \frac{\cos(d_{23})\cos(t) - \cos(d_{12})}{\sin(d_{12})\sin(d_{23})\sin(t)\sin(\alpha_2)}.$$

By the same computation (or a symmetry argument):

$$d\alpha_3(U) = \frac{\cos(d_{01})\cos(t) - \cos(d_{03})}{\sin(d_{01})\sin(d_{03})\sin(t)\sin(\alpha_0)} + \frac{\cos(d_{12})\cos(t) - \cos(d_{23})}{\sin(d_{12})\sin(d_{23})\sin(t)\sin(\alpha_2)}.$$

Proof of Theorem B_S. Let $p = (v_1, \dots, v_n)$ be a convex spherical polygon. Recall that, by Definition 1.2, for any first-order deformation U of p , defined up to the Killing fields, we have:

$$b(U) = \sum_{i=0}^n d\alpha_i(U')dv_i(U'),$$

where U' is any representative of U , i.e. any first-order deformation of p corresponding to U under the quotient by the trivial deformations.

So b is a quadratic form on $T_p\mathcal{P}_S(l)$, where $l = (l_1, \dots, l_n)$ is the n -uple of the edge lengths of p . It is natural to define a bilinear form associated to b , which we call b_2 ; it is defined as follows: if U'_1 and U'_2 are two first-order deformations of p , then:

$$b_2(U'_1, U'_2) := \frac{1}{2} \left(\sum_{i=1}^n d\alpha_i(U'_1)dv_i(U'_2) + \sum_{i=1}^n d\alpha_i(U'_2)dv_i(U'_1) \right).$$

As for b , an important point is that if one adds a Killing field to U'_1 or U'_2 , the result does not change. Moreover, it will be useful below to note that each of the two sums in the definition of b_2 is invariant under this transformation. Indeed, if $Y_1, Y_2 \in \mathbb{R}^3$ are two vectors, let V_1, V_2 be the trivial deformations defined by $dv_i(V_1) = Y_1 \times v_i, dv_i(V_2) = Y_2 \times v_i$; then, for each $i \in \{1, \dots, n\}$, $d\alpha_i(U'_1 + V_1) = d\alpha_i(U'_1)$, and:

$$\begin{aligned} \sum_{i=1}^n d\alpha_i(U'_1 + V_1)dv_i(U'_2 + V_2) &= \sum_{i=1}^n d\alpha_i(U'_1)(dv_i(U'_2) + Y_2 \times v_i) \\ &= \sum_{i=1}^n d\alpha_i(U'_1)dv_i(U'_2) + Y_2 \times \sum_{i=1}^n d\alpha_i(U'_1)v_i \\ &= \sum_{i=1}^n d\alpha_i(U'_1)dv_i(U'_2), \end{aligned}$$

and the same computation can be applied to the second sum in the definition of b_2 .

Lemma 3.1. *Let $p = (v_1, \dots, v_n = v_0)$ be a polygon such that no 3 vertices are collinear. Let U be an isometric first-order deformation of p , such that v_1 and v_2 are fixed. There exists a unique decomposition:*

$$U = U_2 + U_3 + \dots + U_{n-2}$$

such that, for all $i \in \{2, \dots, n-2\}$, U_i vanishes on $v_1, v_2, \dots, v_i$, and acts on $v_{i+1}, \dots, v_{n-1}, v_n = v_0$ as a rigid motion.

Proof. We define a sequence of isometric first-order deformations of p recursively, as follows. First note that, since v_0, v_1 and v_3 are not collinear, there is a unique first-order isometric deformation V_2 of the quadrilateral (v_0, v_1, v_2, v_3) such that v_1 and v_2 are fixed, and that the first-order displacement of v_3 is the same for V_2 and for U . Then define a first-order deformation U_2 as follows:

- v_1 and v_2 are fixed.
- v_3 moves under U_2 as under V_2 , i.e. as under U .
- The restriction of U_2 to $v_3, v_4, \dots, v_n = v_0$ is equal to the restriction to those vertices of a Killing field, i.e. the corresponding part of p moves in a “rigid” way.

It is possible to define such a deformation because, at first order, the distance between v_3 and v_0 does not vary under the deformation V_2 .

Now consider the first-order deformation $U - U_2$ of p . It is isometric — as the difference of two isometric deformations — and it vanishes at v_1, v_2 and v_3 . This shows, because v_1, v_4 and v_0 are not collinear, that there exists a unique isometric first-order deformation V_3 of the quadrilateral (v_1, v_3, v_4, v_0) which vanishes at v_1 and v_3 and acts on v_4 as $U - U_2$. Let U_3 be the unique first-order deformation of p which vanishes at v_1, v_2 and v_3 , and which acts on $v_4, \dots, v_{n-1}, v_0$ as a Killing field (i.e. “rigidly”).

Now $U - U_2 - U_3$ is an isometric first-order deformation of p which vanishes at v_1, v_2, v_3 and v_4 , and we can iterate this construction until we obtain a decomposition:

$$U = U_2 + \dots + U_{n-2} ,$$

which has the property described in the lemma. $\square$

Consider now a first-order deformation U of p . Since p is convex, it is non-degenerate in the sense of Lemma 3.1. Applying this lemma yields a decomposition $U = U_2 + \dots + U_{n-2}$ such that, for all $i \in \{2, \dots, n-2\}$, U_i vanishes on $v_1, v_2, \dots, v_i$, and acts on $v_{i+1}, \dots, v_{n-1}, v_n = v_0$ as a rigid motion.

Lemma 3.2. *For each $i, j \in \{2, \dots, n-2\}$ with $i \neq j$, $\langle b_2(U_i, U_j), v_1 \rangle = 0$. Therefore:*

$$\langle b(U), v_1 \rangle = \left\langle \sum_{i=1}^n b(U_i), v_1 \right\rangle .$$

Proof. Suppose (without loss of generality) that $i < j$. For each $i \in \{2, \dots, n-2\}$, we add a Killing field to U_i to obtain a first-order deformation U'_i which acts on $v_1, v_2, \dots, v_i$ as a rigid motion, and vanishes on $v_{i+1}, \dots, v_{n-1}, v_n = v_0$. It follows from the remark above that:

$$b_2(U_i, U_j) = \frac{1}{2} \left(\sum_{k=1}^n d\alpha_k(U_i) dv_k(U_j) + \sum_{i=1}^n d\alpha_k(U'_j) dv_k(U'_i) \right) .$$

Note that $d\alpha_k(U_i)$ vanishes for all values of k except 0, 1, i and $i+1$. But $dv_k(U_j)$ vanishes, by definition of U_j , for $k = 1, \dots, j$. Symmetrically, $d\alpha_k(U'_j)$ vanishes except for $k = 0, 1, j, j+1$, while $dv_k(U'_i)$ vanishes for $k = i+1, \dots, n$, so that:

$$b_2(U_i, U_j) = \frac{1}{2} (d\alpha_0(U_i) dv_0(U_j) + d\alpha_1(U'_j) dv_1(U'_i)) .$$

Since U_j is an isometric first-order deformation, we have:

$$0 = d\langle v_0, v_1 \rangle(U_j) = \langle dv_0(U_j), v_1 \rangle + \langle v_0, dv_1(U_j) \rangle .$$

But $dv_1(U_j) = 0$ by definition of U_j , so that $\langle dv_0(U_j), v_1 \rangle = 0$. Since $dv_1(U'_i) \in T_{v_1} S^2$, $\langle v_1, dv_1(U'_i) \rangle = 0$, and it follows that $\langle b_2(U_i, U_j), v_1 \rangle = 0$. $\square$

The value of $\langle b(U_i), v_1 \rangle$ is given by the next lemma.

Lemma 3.3. *Let $q := (v_1, v_i, v_{i+1}, v_0)$, and let $\alpha'_1, \alpha'_i, \alpha'_{i+1}, \alpha'_0$ be its angles. Let V be the first-order isometric deformation of q which vanishes on v_1 and v_i and under which the distance between v_1 and v_{i+1} varies at speed 1. Then, if $d_{j,k} := d(v_j, v_k)$, we have:*

$$\langle v_1, d\alpha'_{i+1}(V)dv_{i+1}(V) + d\alpha'_0(V)dv_0(V) \rangle = \frac{\sin^2(d_{1,i+1}) \sin(\alpha'_1)}{\sin(d_{0,i+1}) \sin(d_{i,i+1}) \sin(\alpha'_0) \sin(\alpha'_i)} .$$

Proof. We have already noted that $\langle v_1, dv_0(V) \rangle = 0$ because V does not change, at first order, the distance between v_0 and v_1 . So the only non-vanishing term is the one involving $\langle v_1, dv_{i+1}(V) \rangle$, and:

$$\langle v_1, dv_{i+1}(V) \rangle = d\langle v_1, v_{i+1} \rangle(V) = d\cos(d_{1,i+1})(V) = -\sin(d_{1,i+1}) .$$

But $d\alpha'_{i+1}(V)$ is given by equation (1); we now have slightly different notations, U is replaced by V , the indices 2 (resp. 3) by i (resp. $i+1$). So:

$$d\alpha'_{i+1}(V) = \frac{\cos(d_{0,1}) \cos(t) - \cos(d_{0,i+1})}{\sin(d_{0,1}) \sin(d_{0,i+1}) \sin(t) \sin(\alpha'_0)} + \frac{\cos(d_{1,i}) \cos(t) - \cos(d_{i,i+1})}{\sin(d_{1,i}) \sin(d_{i,i+1}) \sin(t) \sin(\alpha'_i)} ,$$

where $t := d_{1,i+1}$. Following the notations of Figure 3, we call β'_1 and β'_{i+1} the angles at v_1 and v_{i+1} , respectively, of the triangle (v_0, v_1, v_{i+1}) , and we call γ'_1 and γ'_{i+1} the angles at v_1 and v_{i+1} , respectively, of the triangle (v_1, v_i, v_{i+1}) . Using equation (1), we get:

$$\begin{aligned} d\alpha'_{i+1}(V) &= -\frac{\cos(\beta'_1) \sin(d_{0,1}) \sin(t)}{\sin(d_{0,1}) \sin(d_{0,i+1}) \sin(t) \sin(\alpha'_0)} - \frac{\cos(\gamma'_1) \sin(d_{1,i}) \sin(t)}{\sin(d_{1,i}) \sin(d_{i,i+1}) \sin(t) \sin(\alpha'_i)} \\ &= -\sin(t) \left(\frac{\cos(\beta'_1)}{\sin(d_{0,i+1}) \sin(\alpha'_0) \sin(t)} + \frac{\cos(\gamma'_1)}{\sin(d_{i,i+1}) \sin(\alpha'_i) \sin(t)} \right) \\ &= -\frac{\sin(t)}{\sin(d_{0,i+1}) \sin(d_{i,i+1}) \sin(\alpha'_0) \sin(\alpha'_i)} \times \\ &\times \left(\cos(\beta'_1) \frac{\sin(d_{i,i+1}) \sin(\alpha'_i)}{\sin(t)} + \cos(\gamma'_1) \frac{\sin(d_{0,i+1}) \sin(\alpha'_0)}{\sin(t)} \right) . \end{aligned}$$

But, by the sine law for spherical triangles:

$$\frac{\sin(\alpha'_i)}{\sin(t)} = \frac{\sin(\gamma'_1)}{\sin(d_{i,i+1})} , \quad \frac{\sin(\alpha'_0)}{\sin(t)} = \frac{\sin(\beta'_1)}{\sin(d_{0,i+1})} ,$$

so that:

$$\begin{aligned} d\alpha'_{i+1}(V) &= -\sin(t) \frac{\cos(\beta'_1) \sin(\gamma'_1) + \cos(\gamma'_1) \sin(\beta'_1)}{\sin(d_{0,i+1}) \sin(d_{i,i+1}) \sin(\alpha'_0) \sin(\alpha'_i)} \\ &= -\sin(t) \frac{\sin(\alpha'_1)}{\sin(d_{0,i+1}) \sin(d_{i,i+1}) \sin(\alpha'_0) \sin(\alpha'_i)} , \end{aligned}$$

and the result follows. $\square$

It follows directly from Lemma 3.2 and from Lemma 3.3 that $\langle b(U), v_1 \rangle$ is a sum of positive terms, so it is positive. By symmetry the same holds for all the other vertices. Since $b(U)$ has positive scalar product with all the vertices of p , it is contained in the positive cone over the interior of the dual polygon p^* , and this proves Theorem B_S .

Proof of Theorem B_H . The same proof applies in the hyperbolic case, with some small differences in the computations. We only state the hyperbolic analog of Lemma 3.3.

Lemma 3.4. *Let $q := (v_1, v_i, v_{i+1}, v_0)$, and let $\alpha'_1, \alpha'_i, \alpha'_{i+1}, \alpha'_0$ be its angles. Let V be the first-order isometric deformation of q which vanishes on v_1 and v_i and under which the distance between v_1 and v_{i+1} varies at speed 1. Then, if $d_{j,k} := d(v_j, v_k)$, we have:*

$$\langle v_1, d\alpha'_{i+1}(V)dv_{i+1}(V) + d\alpha'_0(V)dv_0(V) \rangle = \frac{\sinh^2(d_{1,i+1}) \sin(\alpha'_1)}{\sinh(d_{0,i+1}) \sinh(d_{i,i+1}) \sin(\alpha'_0) \sin(\alpha'_i)} .$$

The proof is based on some computations which are quite parallel to those made above for first-order deformations of spherical quadrilaterals, but now for hyperbolic quadrilaterals. We use again the notations apparent in Figure 3. The basic triangle formula is now:

$$\cosh(t) = \cosh(d_{01}) \cosh(d_{03}) - \cos(\alpha_0) \sinh(d_{01}) \sinh(d_{03}) ,$$

from which it follows that, in first-order deformation U such that t varies at speed 1, we have:

$$d\alpha_0(U) = \frac{\sinh(t)}{\sin(\alpha_0) \sinh(d_{01}) \sinh(d_{03})} ,$$

and the same computation shows that:

$$d\alpha_2(U) = \frac{\sinh(t)}{\sin(\alpha_2) \sinh(d_{12}) \sinh(d_{23})} .$$

Moreover, equation (1) yields:

$$\cos(\beta_1) = \frac{-\cosh(d_{03}) + \cosh(d_{01}) \cosh(t)}{\sinh(d_{01}) \sinh(t)} ,$$

so that:

$$\begin{aligned} -\sin(\beta_1)d\beta_1(U) &= \frac{\cosh(d_{01}) \sinh^2(t) - (-\cosh(d_{03}) + \cosh(d_{01}) \cosh(t)) \cosh(t)}{\sinh(d_{01}) \sinh^2(t)} \\ &= \frac{-\cosh(d_{01}) + \cosh(d_{03}) \cosh(t)}{\sinh(d_{01}) \sinh^2(t)} . \end{aligned}$$

But:

$$\frac{\sin(\beta_1)}{\sinh(d_{03})} = \frac{\sin(\alpha_0)}{\sinh(t)} ,$$

so that:

$$d\beta_1(U) = \frac{-\cosh(d_{03}) \cosh(t) + \cosh(d_{01})}{\sinh(d_{01}) \sinh(d_{03}) \sinh(t) \sin(\alpha_0)} .$$

By symmetry:

$$d\gamma_1(U) = \frac{-\cosh(d_{23}) \cosh(t) + \cosh(d_{12})}{\sinh(d_{12}) \sinh(d_{23}) \sinh(t) \sin(\alpha_2)} ,$$

and, taking, the sum:

$$d\alpha_1(U) = \frac{-\cosh(d_{03}) \cosh(t) + \cosh(d_{01})}{\sinh(d_{01}) \sinh(d_{03}) \sinh(t) \sin(\alpha_0)} + \frac{-\cosh(d_{23}) \cosh(t) + \cosh(d_{12})}{\sinh(d_{12}) \sinh(d_{23}) \sinh(t) \sin(\alpha_2)} .$$

By the same computation (or a symmetry argument):

$$d\alpha_3(U) = \frac{-\cosh(d_{01}) \cosh(t) + \cosh(d_{03})}{\sinh(d_{01}) \sinh(d_{03}) \sinh(t) \sin(\alpha_0)} + \frac{-\cosh(d_{12}) \cosh(t) + \cosh(d_{23})}{\sinh(d_{12}) \sinh(d_{23}) \sinh(t) \sin(\alpha_2)} .$$

The end of the proof then proceeds as in the spherical case, we leave the details to the interested reader.

The submanifold of angles for given edge lengths. One possible way to interpret geometrically the positivity property of b for spherical polygons is in terms of a kind of convexity property of a submanifold of codimension 3 in $\mathbb{R}^n$, defined as the set of possible angles of convex polygons with given edge lengths. This submanifold is often smooth, and when it has singularities, they are located at precise points. Its tangent space at each point is given by Theorem A_S , while Theorem B_S indicates that its second fundamental form is always on “one side”, as explained below.

We consider here a spherical polygon p with n vertices, and call $l = (l_1, \dots, l_n)$ the family of its edge length. Then we call $\mathcal{A}_S(l)$ the space of n -uples $(\alpha_1, \dots, \alpha_n)$ such that there exists a polygon $p' \in \mathcal{P}_S(l)$ with angles equal to the α_i . This defines an map Φ_l from $\mathcal{P}_S(l)$ into $\mathbb{R}^n$, which is with image $\mathcal{A}_S(l)$, because a spherical polygon is entirely defined – up to global isometries in S^2 – by its edge lengths and its angles.

Theorem A_S , along with the remarks at the beginning of section 2, show that $\mathcal{A}_S(l)$ is locally a smooth submanifold of codimension 3 of $(\mathbb{R}/2\pi\mathbb{Z})^n$, except when all the vertices of the polygon lie on a spherical geodesic. Clearly this is possible only when there exists $(\epsilon_1, \dots, \epsilon_n) \in \{-1, 1\}$ and $k \in \mathbb{Z}$ such that:

$$\sum_{i=1}^n \epsilon_i l_i = 2k\pi .$$

Then the singular points of $\mathcal{A}_S(l)$ can happen only at the points of $(\mathbb{R}/2\pi\mathbb{Z})^n$ which have all their coordinates equal to 0 or π .

Let $q \in P_S(l)$, with vertices $v_1, \dots, v_n$. Let $\alpha_1, \dots, \alpha_n$ be the angles of q , we suppose that they are not all equal to 0 or π , so that $\mathcal{A}_S(l)$ is locally smooth in the neighborhood of $\alpha := (\alpha_1, \dots, \alpha_n)$. The tangent space of $\mathcal{A}_S(l)$ is described by Theorem A_S . More precisely, there is a natural isomorphism between $\mathbb{R}^3$ and the normal space of $\mathcal{A}_S(l)$ at α , defined as follows:

$$\forall w \in \mathbb{R}^3, \Psi_q(w) := (\langle v_1, w \rangle, \dots, \langle v_n, w \rangle) .$$

Theorem A_S shows that $\Psi_q(\mathbb{R}^3)$ is the orthogonal to the tangent space of $\mathcal{A}_S(l)$ at α , i.e. the normal space of $\mathcal{A}_S(l)$ at α .

Now b appears naturally as the second fundamental form of $\mathcal{A}_S(l)$.

Remark 3.5. Let $U \in T_q P_S(l)$ be a first-order isometric deformation of q , and let $w \in \mathbb{R}^3$. then:

$$\langle \mathcal{I}(d\Phi_l(U), d\Phi_l(U)), \Psi_q(w) \rangle = -\langle b(U), w \rangle ,$$

where $\mathcal{I}$ is the second fundamental form of $\mathcal{A}_S(l)$.

Proof. Let $w \in \mathbb{R}^3$. Consider any extension of U as a vector field tangent to $P_S(l)$ in the neighborhood of q , and call ∇ the flat connection on $\mathbb{R}^n$. By definition of the second fundamental form of $\mathcal{A}_S(l)$, we have:

$$\begin{aligned} \langle \mathcal{I}(d\Phi_l(U), d\Phi_l(U)), \Psi_q(w) \rangle &= -\langle \nabla_{d\Phi_l(U)} \Psi_q(w), d\Phi_l(U) \rangle \\ &= -\langle (\langle dv_i(U), w \rangle)_{i=1}^n, (d\alpha_i(U))_{i=1}^n \rangle \\ &= -\sum_{i=1}^n \langle d\alpha_i(U) dv_i(U), w \rangle \\ &= -\langle b(U), w \rangle . \end{aligned}$$

□

There is a direct translation of Theorem B_S in this context, indicating a kind of convexity property of $\mathcal{A}_S(l)$ at the points which are the images of convex polygons.

Proposition 3.6. Suppose that q is convex. Let Q^* be the polyhedral cone in $N_{\Phi_l(q)} \mathcal{A}_S(l)$ which is dual to the cone $\Psi_q((\mathbb{R}_+^*)^n)$, for the metric induced on $N_q \mathcal{A}_S(l)$ by the metric on $\mathbb{R}^n$. Then:

$$\forall V \in T_{\Psi_l(q)} \mathcal{A}_S(l), \mathcal{I}(V, V) \in -Q^* .$$

Proof. By the previous remark and Theorem B_S:

$$\forall U \in T_q P_S(l), \forall w \in \mathbb{R}_+^* q, \langle \mathcal{H}(d\Phi_l(U), d\Phi_l(U)), \Psi_q(w) \rangle < 0 ,$$

so that:

$$\forall V \in T_{\Phi_l(q)} \mathcal{A}_S(l), \forall W \in \Psi_q(\mathbb{R}_+^* q), \langle \mathcal{H}(V, V), W \rangle < 0 ,$$

and the result follows. $\square$

The same considerations hold also for hyperbolic polygons, based on Theorem A_H for the description of the tangent space, and on Theorem B_H for the convexity property of the submanifold of the angles of polygons with given edge lengths.

In the Euclidean case, one can again consider the space $\mathcal{A}_E(l)$ of angles of polygons with edge lengths given by l . Then $\mathcal{A}_E(l)$ is a codimension 2 submanifold of the hyperplane of equation: $\sum_{i=1}^n \alpha_i = 0$. However it does not appear to have any obvious “convexity” property.

4 Isoperimetric problems for spherical or hyperbolic polygons

The theorems given above – Theorem A_S and Theorem B_S for spherical polygons, Theorem A_H and Theorem B_H for hyperbolic polygons – have very simple applications to isoperimetric questions in the two settings. The spherical result was apparently discovered by Steiner [Ste42], and is analogous to an Euclidean statement which was known earlier (see e.g. [Sie02] for recent progress on related but more elaborate Euclidean statements). The proofs are strongly related to Theorems A_S and B_S (resp. A_H and B_H).

Critical points of the area functional. First, Theorem A_S and Theorem A_H lead directly to a description of the critical points of the area function over the space of polygons with given edge lengths.

Lemma 4.1. *Let p be a spherical polygon, with edge lengths given by $l = (l_1, \dots, l_n)$. Suppose that the vertices of p are not all on a geodesic. Then p is a critical point of the area in $\mathcal{P}_S(l)$ if and only if all vertices of p are on a circle.*

Lemma 4.2. *Let p be a hyperbolic polygon, with edge lengths given by $l = (l_1, \dots, l_n)$. Suppose that the vertices of p are not all on a geodesic. Then p is a critical point of the area in $\mathcal{P}_H(l)$ if and only if all vertices of p are either on a circle, on a horocycle, or on a connected component of the set of points at fixed distance from a geodesic.*

Note that the polygons are not required to be convex. It follows from the results given below that, for convex polygons, the critical points are actually maxima, but it is probably not the case for non-convex polygons. Both lemmas are proved together.

Proof. By the Gauss-Bonnet formula, p is a critical point of the area functional if and only if, for any first-order isometric deformation of p , the sum of the angles of p remains constant. By Theorem A_S (resp. Theorem A_H) this happens if and only if:

$$\forall \dot{\alpha}_1, \dots, \dot{\alpha}_n \in \mathbb{R}, \left(\sum_{i=1}^n \dot{\alpha}_i v_i = 0 \Rightarrow \sum_{i=1}^n \dot{\alpha}_i = 0 \right) .$$

By an elementary linear algebra argument using the transposed equation, this is true if and only if there exists a vector $u \in \mathbb{R}^3$ (resp. $\mathbb{R}_1^3$) such that:

$$\forall i \in \{1, \dots, n\}, \langle u, v_i \rangle = 1 ,$$

which holds if and only if the v_i are on an affine plane in $\mathbb{R}^3$ (resp. $\mathbb{R}_1^3$) not containing the origin. The result follows, because the intersection of S^2 with an affine plane in $\mathbb{R}^3$ is a circle (or a point) while the intersection of H^2 with a plane in $\mathbb{R}_1^3$ is either a circle, a horocycle, or a connected component of the set of points at fixed distance from a geodesic. $\square$

Uniqueness of critical polygons. In light of the previous paragraph, it is interesting to remark that, for a given set of edge lengths, there is at most one convex polygon which is a critical point of the area. We first consider spherical polygons, for which the result has been well-known for many years.

Lemma 4.3. *Let $l_1, \dots, l_n \in \mathbb{R}_+^*$. There is at most one convex polygon $p \in \mathcal{P}_S^c(l)$ which has its vertices on a circle.*

Proof. We suppose that l_1 is the largest of the l_i .

Suppose that the vertices of p are on a circle of radius l and center x_0 in S^2 . Then the edge of p of length l_i is “seen” from x_0 under an angle θ_i , and the sine law for spherical triangles shows that:

$$\sin(\theta_i/2) = \frac{\sin(l_i/2)}{\sin(l)}.$$

Setting $S_i := \sin(l_i/2)$ and $S := \sin(l)$, we have again two possibilities:

- Either x_0 is in the interior of p , and then:

$$\sum_{i=1}^n \arcsin(S_i/S) = \pi.$$

- Or x_0 is not in the interior of p , and then:

$$\sum_{i=2}^n \arcsin(S_i/S) = \arcsin(S_1/S).$$

Moreover, in both cases, we have:

$$\sum_{i=2}^n \arcsin(S_i/S) < \pi,$$

and:

$$S_1/S = \sin\left(\sum_{i=2}^n \arcsin(S_i/S)\right).$$

Set:

$$F(s) := s \sin\left(\sum_{i=2}^n \arcsin\left(\frac{S_i}{s}\right)\right).$$

Then:

$$\begin{aligned} F'(s) &= \sin\left(\sum_{i=2}^n \arcsin\left(\frac{S_i}{s}\right)\right) + s \cos\left(\sum_{i=2}^n \arcsin\left(\frac{S_i}{s}\right)\right) \left(\sum_{i=2}^n \frac{-S_i/s^2}{\sqrt{1 - S_i^2/s^2}}\right) \\ &= \cos\left(\sum_{i=2}^n \arcsin\left(\frac{S_i}{s}\right)\right) \left[\tan\left(\sum_{i=2}^n \arcsin\left(\frac{S_i}{s}\right)\right) - \sum_{i=2}^n \tan\left(\arcsin\left(\frac{S_i}{s}\right)\right)\right]. \end{aligned}$$

When $\sum_{i=2}^n \arcsin(S_i/s) \in [\pi/2, \pi)$, both terms in the first equation are non-positive, and it follows that $F'(s)$ is positive. When $\sum_{i=2}^n \arcsin(S_i/s) \in (0, \pi/2)$, the second equation shows the same result, because the cosine is positive, while $\tan$ is convex on $(0, \pi/2)$, so that the second product is also positive. Therefore, F is increasing, and it follows that there is at most one s such that $F(s) = S_1$, and therefore at most one possible value of $S = \sin(l)$.

But, given l , the angles α_i are uniquely determined by the l_i , so that there is at most one polygon with vertices on a circle and with edge lengths equal to the l_i . $\square$

We now consider the hyperbolic case. This is a step in the proof of Theorem 1.4.

Lemma 4.4. *Let $l_1, \dots, l_n \in \mathbb{R}_+^*$. There is at most one convex polygon $p \in \mathcal{P}_H^c(l)$ which has its vertices either on a circle, on a horocycle, or on a connected component of the set of points at constant distance from a geodesic.*

Proof. We suppose again that l_1 is the largest of the l_i .

Suppose that the vertices of p are on a circle of radius l and center x_0 in H^2 . Then the edge of p of length l_i is “seen” from x_0 under an angle θ_i , and the sine law for hyperbolic triangles shows that:

$$\sin(\theta_i/2) = \frac{\sinh(l_i/2)}{\sinh(l)} .$$

Setting $S_i := \sinh(l_i/2)$ and $S := \sinh(l)$, we have again two possibilities:

- Either x_0 is in the interior of p , and then:

$$\sum_{i=1}^n \arcsin(S_i/S) = \pi .$$

- Or x_0 is not in the interior of p , and then:

$$\sum_{i=2}^n \arcsin(S_i/S) = \arcsin(S_1/S) .$$

Moreover, in both cases, we have:

$$S_1/S = \sin \left(\sum_{i=2}^n \arcsin(S_i/S) \right)$$

and, since $\sin$ is concave on $[0, \pi]$:

$$S_1/S < \sum_{i=2}^n \sin(\arcsin(S_i/S)) = \sum_{i=2}^n S_i/S ,$$

so that $S_1 < \sum_{i=2}^n S_i$.

Suppose now that the vertices of p are on a horocycle h . An elementary argument in hyperbolic geometry shows that the distance, along h , between the vertices of an edge of length l_i is equal to $2 \sinh(l_i/2)$.

In the projective model of H^2 , the horocycle h appears as an ellipse, tangent to the boundary at infinity at a point x_0 . In this model, p appears as a convex polygon, with vertices on this ellipse. There is exactly one edge e of p such that x_0 is on one side of e , while all the other edges of p are on the other side. Clearly, e has to be the edge of p of largest lengths, and, in this case:

$$S_1 = \sum_{i=2}^n S_i .$$

Finally, suppose that all vertices of p are on a connected component c of the set of points at distance l from a geodesic g_0 . Let $e_1, \dots, e_n$ be the edges of p ; the sine formula for hyperbolic triangles shows that, if a_i is the distance between the orthogonal projections on g_0 of e_i , then:

$$\sinh(a_i/2) = \frac{\sinh(l_i/2)}{\sinh(l)} .$$

In the projective model of H^2 , the set of points at distance l from g_0 appears as an ellipse, tangent to the boundary at infinity of H^2 at the endpoints of g_0 , and p appears as an ellipse (see Figure 4). By




Figure 2: The set of points at constant distance from a geodesic.

convexity, there is one edge e of p such that g_0 is on one side, and all the other edges of p on the other. This edge has maximal length, and:

$$a_1 = \sum_{i=2}^n a_i ,$$

so that:

$$\operatorname{argsinh}(S_1/S) = \sum_{i=2}^n \operatorname{argsinh}(S_i/S) .$$

Then, taking the sinh, we have, because sinh is convex on $(0, \infty)$:

$$S_1/S = \sinh \left(\sum_{i=2}^n \operatorname{argsinh}(S_i/S) \right) > \sum_{i=2}^n S_i/S ,$$

so that $S_1 > \sum_{i=2}^n S_i$.

Considering those three cases, it is clear that the l_i determine whether there can exist a convex polygon p with edge lengths equal to the l_i and vertices on a circle, a horocycle, or a connected component of the set of points at constant distance from a geodesic.

1. If $S_1 < \sum_{i=2}^n S_i$, then the only possibility is that the vertices of p are on a circle.
2. If $S_1 = \sum_{i=2}^n S_i$, then the vertices of p can only be on a horocycle.
3. If $S_1 > \sum_{i=2}^n S_i$, the vertices of p can only be on a connected component of the set of points at constant distance from a geodesic.

In the first case, we proceed as in the proof of Lemma 4.3 above, and set:

$$F(s) := s \sin \left(\sum_{i=2}^n \arcsin \left(\frac{S_i}{s} \right) \right) .$$

Either equation (1) or equation (1) is satisfied; in both cases, $F(S) = S_1$. However the same proof as for Lemma 4.3 shows that F is strictly increasing, so that there is at most one possible value of S . As in the spherical case, it is easy to check that a polygon with vertices on a circle is uniquely determined (up to global isometries) by its edge lengths and the radius of the circle; this finishes the proof case (1).

In the second case, the polygon p is clearly uniquely determined (up to global isometries) by the distances between its vertices on the horocycle, so the statement in the lemma holds.

In case (3), it is necessary to modify slightly the argument used in the spherical case. We set:

$$G(s) := s \sinh \left(\sum_{i=2}^n \operatorname{argsinh} \left(\frac{S_i}{s} \right) \right) .$$

Now:

$$\begin{aligned} G'(s) &= \sinh \left(\sum_{i=2}^n \operatorname{argsinh} \left(\frac{S_i}{s} \right) \right) + s \cosh \left(\sum_{i=2}^n \operatorname{argsinh} \left(\frac{S_i}{s} \right) \right) \left(\sum_{i=2}^n \frac{-S_i/s^2}{\sqrt{1 + S_i^2/s^2}} \right) \\ &= \cosh \left(\sum_{i=2}^n \operatorname{argsinh} \left(\frac{S_i}{s} \right) \right) \left[\tanh \left(\sum_{i=2}^n \operatorname{argsinh} \left(\frac{S_i}{s} \right) \right) - \sum_{i=2}^n \tanh \left(\operatorname{argsinh} \left(\frac{S_i}{s} \right) \right) \right] . \end{aligned}$$

But the first factor in the right-hand term is positive, while the second factor is negative because $\tanh$ is concave on $(0, \infty)$. So G is strictly decreasing, and there is at most one value of s such that $G(s) = S_1$. It follows that there is at most one polygon with edge lengths equal to the l_i and with vertices on a connected component of the set of points at constant distance from a geodesic. $\square$

Boundary behavior of the area functional. Another, simple element in the proof of Theorem 1.4 is that, on the boundary of the space $\mathcal{P}_S^c(l)$ (resp. $\mathcal{P}_H^c(l)$), the interior derivative of the area is positive.

Lemma 4.5. *Let $p \in \partial \mathcal{P}_S^c(l)$ (resp. $p \in \partial \mathcal{P}_H^c(l)$) be a polygon with exactly one angle equal to π . Then the interior normal derivative of the area at p is positive.*

Proof. Let v be the vertex of p at which the angle is equal to π . Consider the isometric first-order deformation U of p under which v moves toward the exterior of p , at speed 1, along the orthogonal to both the edges of p at v . Clearly, U is towards the interior of $\mathcal{P}_S^c(l)$ (resp. of $\mathcal{P}_H^c(l)$) and the first-order variation of the area under U is positive. The result follows. $\square$

Proof of Theorem 1.4. By Lemma 4.5, the area has at least one local maximum in each connected component of $\mathcal{P}_S^c(l)$ (resp. of $\mathcal{P}_H^c(l)$). By Lemma 4.2, each critical point of the area is a polygon with vertices on a circle, a horocycle, or a connected component of the set of points at constant distance from a geodesic. By Lemma 4.4, there is at most one such critical point. So the area has exactly one critical point in $\mathcal{P}_H^c(l)$, which is a maximum. By the way, it also follows that $\mathcal{P}_H^c(l)$ is connected.

Note that the same argument can be used for spherical polygons, it prove the corresponding result.

The convexity of the area. In addition, Theorem B_S (resp. B_H) shows that the area has a strict concavity property over at least a subset of the space of convex polygon with given edge lengths. We call A the function equal to the area of polygons.

Definition 4.6. *Let p be a convex polygon, with vertices $v_1, \dots, v_n$. Let $\phi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be defined by:*

$$\phi(x) = \sum_{i=1}^n \langle x, v_i \rangle v_i .$$

Suppose that the v_i are not on a geodesic, let C_v be the vertices of p ; we set:

$$C_S(p) := \frac{\phi^{-1}(C_v)}{\|\phi^{-1}(C_v)\|} .$$

Note that, since the v_i are not on a geodesic, ϕ is a self-adjoint, positive definite operator, so it is invertible.

To understand the meaning of $C_S(p)$, we consider first the simplest case, when p is a spherical triangle, with vertices v_1, v_2 and v_3 , which we suppose are not on a geodesic. Then $C_S(p)$ is the barycenter of the

dual polygon p^* . Indeed, let $p^* = (v_1^*, v_2^*, v_3^*)$; then (v_1^*, v_2^*, v_3^*) is the basis of $\mathbb{R}^3$ which is dual to the basis (v_1, v_2, v_3) . By definition, $C_S(p)$ is defined by the equation:

$$\frac{\sum_{i=1}^3 \langle v_i, C_S(p) \rangle v_i}{\left\| \sum_{i=1}^3 \langle v_i, C_S(p) \rangle v_i \right\|} = C_v(p) .$$

Taking the scalar product with v_j^* , we obtain that $C_S(p)$ is characterized by the existence of a $\lambda > 0$ such that, for all $j \in \{1, 2, 3\}$:

$$\langle C_S(p), v_j \rangle = \lambda \langle C_v(p), v_j^* \rangle .$$

The action of the duality on this equation shows that $C_S(p)$ is the barycenter of the polygon $p^* = (v_1^*, v_2^*, v_3^*)$.

This example can be extended by taking a triangle with integer “weights”, i.e. we consider a polygon with $k_1 + k_2 + k_3$ vertices, k_1 of them being “collapsed” on a point v_1 , k_2 on v_2 , and k_3 on v_3 . The argument given above carries over to this case, and, again in this case, the point $C_S(p)$ which is obtained is the barycenter of the triangle dual to (v_1, v_2, v_3) ; in other terms, the weights k_1, k_2, k_3 have no influence. This illustrates an interesting “stability” property of $C_S(p)$.

The concavity of the area. Theorem 1.4 is related to the fact that, over at least part of $\mathcal{P}_H^c(l)$ (resp. $\mathcal{P}_S^c(l)$), the area is strictly concave.

Lemma 4.7. *Let p be a convex polygon in S^2 (resp. H^2) with edge lengths given by $l = (l_1, \dots, l_n)$. Suppose that $C_S(p)$ is contained in the interior of p . Then the restriction of A to $\mathcal{P}_S^c(l)$ is strictly concave at p for the metric induced by the immersion of $\mathcal{P}_S(l)$ in $[0, \pi]^n$ defined by sending a polygon to its angles.*

Proof. First note that A has an extension as a linear function over $[0, \pi]^n$ since, by the Gauss-Bonnet formula, $A = 2\pi - \sum_{i=1}^n (\pi - \alpha_i)$ for spherical polygons, while $A = -2\pi + \sum_{i=1}^n (\pi - \alpha_i)$ for hyperbolic polygons. We still call this function A , then $dA = (1, \dots, 1)$ for spherical polygons, while $dA = -(1, \dots, 1)$ for hyperbolic polygons.

Let $\text{Hess}(A)$ be the Hessian of A over $\Phi_l(\mathcal{P}_S(l))$. Let U be a first-order isometric deformation of p , then:

$$\text{Hess}(A)(U, U) = dA(\text{II}(d\Phi_l(U), d\Phi_l(U))) = \langle \text{II}(d\Phi_l(U), d\Phi_l(U)), I \rangle ,$$

where II is the second fundamental form of $\Phi_l(\mathcal{P}_S^c(l))$ and I is the vector $(1, \dots, 1)$. Since $\text{II}(d\Phi_l(U), d\Phi_l(U))$ is orthogonal to $\Phi_l(\mathcal{P}_S^c(l))$:

$$\text{Hess}(A)(U, U) = \langle \text{II}(d\Phi_l(U), d\Phi_l(U)), I_N \rangle ,$$

where I_N is the orthogonal projection of I on the normal space of $\Phi_l(\mathcal{P}_S^c(l))$ at $\Phi_l(p)$. By definition of Ψ_p , $I_N = \Psi_p(w)$, where $w \in \mathbb{R}^3$ is characterized by the fact that $\Psi_p(w) = I_N$, i.e. that $\Psi_p(w) - I \in T_{\Phi_l(p)}\Phi_l(\mathcal{P}_S^c(l))$, which translates as:

$$\sum_{i=1}^n (\langle w, v_i \rangle - 1) v_i = 0 ,$$

so that:

$$\sum_{i=1}^n \langle w, v_i \rangle v_i = \sum_{i=1}^n v_i .$$

This shows that $w = \lambda C_S(p)$, for some $\lambda > 0$.

Applying Remark 3.5 then leads to:

$$\text{Hess}(A)(U, U) = -\lambda \langle b(U), C_S(p) \rangle ,$$

so that the Hessian of A is negative definite at p .

The proof in the spherical case is almost the same, with some differences in the signs, and we leave it to the interested reader. $\square$

Although the proof given here only works when $C_S(p)$ is contained in the interior of p , it is not obvious whether the result is valid for all convex polygons – the existence of a unique critical point of the area over $\mathcal{P}_S^c(l)$, which is true in all cases, suggest that it might be the case.

5 Infinitesimal rigidity of Euclidean polyhedra

Infinitesimal rigidity of convex polyhedra. In this section we consider convex polyhedra in $\mathbb{R}^3$. We define a convex polyhedron as the boundary of a compact subset of $\mathbb{R}^3$ which is the intersection of a finite number of half-spaces. Given a polyhedron P , which is the boundary of a compact subset $\Omega \subset \mathbb{R}^3$, where Ω is the intersection of distincts closed half-spaces $H_1, \dots, H_n$ (each of which intersect the interior of Ω , a vertex of P is a point of P which is contained in at least 3 of the boundaries of the H_i . An edge of P is a maximal connected subset of P which is contained in the boundaries of two of the H_i but containing no vertex. A face of P is a connected component of the complement of the vertices and the edges.

The geometry of Euclidean polyhedra has interested geometers for quite a long time [Euc02]. A nice result of Legendre [LegII]² and Cauchy [Cau13] states that convex polyhedra are rigid: if two convex polyhedra have the same combinatorics and the same induced metric on their faces, they are congruent. A related result, first proved by Dehn [Deh16] (which is also a consequence of the ideas of Legendre and Cauchy) is that convex polyhedra are infinitesimally rigid: any first-order deformation of a polyhedron P which acts as a Killing field on each of its faces is the restriction to P of a global Killing field.

From a modern viewpoint, one of the main motivation to study the infinitesimal rigidity of convex polyhedra is that it is the key point in the proof of the following nice theorem.

Theorem 5.1 (Aleksandrov [Ale58]). *The induced metric on any convex polyhedron in $\mathbb{R}^3$ is flat, with conical singularities where the total angle is less than 2π . Conversely, any flat metric on S^2 with conical singularities where the total angle is less than 2π is induced on a unique (up to isometries) convex polyhedron in $\mathbb{R}^3$.*

To prove this theorem, one considers the natural map sending a convex polyhedron to its induced metrics; it is a map between two spaces of the same dimension, and the infinitesimal rigidity statement means that its differential is everywhere an isomorphism, so that it is a local homeomorphism.

We will show here that Theorem C is a consequence of Theorem B_S, thus providing a new proof which is rather natural. It will also serve as an introduction for the rigidity result of the next section, since the proof given there is related but more complicated. The rigidity proof given here is related to a proof of the Cauchy-Legendre rigidity result discovered by Pogorelov [Pog56].

Closed convex polyhedra. Let P be a closed, convex Euclidean polyhedron. Choose a point p_0 in the interior of P . Let u_0 be the function defined on $\mathbb{R}^3$ by: $u_0(p) = d(p_0, p)^2/2$. We consider a first-order deformation of P , i.e. a vector field v on P which acts as a Killing field on each of its faces; we will show that v is trivial, i.e. it acts as a Killing field on P as a whole.

Let e be an edge of P . The first-order deformation v of P determines a first-order variation $\dot{\theta}_e$ of the dihedral angle θ_e of P at e . v also determines a function, which we call $\dot{u}_0$, on P , defined as the first-order variation of the restriction of u_0 to P . By definition of u_0 , we have:

$$\forall p \in P, \quad \dot{u}_0(p) = \langle v(p), p_0 p \rangle .$$

We define a 1-form W on the 1-skeleton of P (the union of its edges) as follows: for each vector u tangent to an edge e , $W(u) = \dot{\theta}_e d\dot{u}_0(u)$.

Consider a parametrization of e at constant speed, say: $p(t) = at + b$, with $a, b \in \mathbb{R}^3$. Then, using (1):

$$\dot{u}_0(p(t))' = d\dot{u}_0(p'(t)) = \langle dv(p'(t)), p_0 p(t) \rangle + \langle v(p(t)), p'(t) \rangle .$$

But $p'(t) = a$ is independent of t , and $dv(p'(t))$ does not depend on t either since v acts on e as a Killing field. So:

$$\frac{d}{dt} \dot{u}_0(p(t))' = 2 \langle dv(p'(t)), p'(t) \rangle .$$

But v acts isometrically on e , so that $\langle dv(p'(t)), p'(t) \rangle = 0$, and it follows that $d\dot{u}_0(p'(t))$ is constant over e , so $\dot{\theta}_e d\dot{u}_0(p'(t))$ is constant over e .

²The contribution of Legendre was recently pointed out by I. Sabitov

Now consider an oriented edge e of P , we call e_- and e_+ its endpoints, and we let W_e be the number $W(p'(t) = \dot{\theta}_e d\dot{u}_0(p'(t)))$, where $p(t)$ is a parametrization of e at speed 1, respecting the orientation. Then we have:

$$\sum_v \sum_{e_- = v} W_e = 0 ,$$

because each non-oriented edge contributes twice, with opposite signs.

Let x be one of the vertices of P . Let $e_1, \dots, e_{n-1}, e_n = e_0$ be the oriented edges of P such that $e_{i,-} = x$, with their natural cyclic order, and let $v_1, \dots, v_n = v_0$ be unit vectors with v_i in the direction of e_i , oriented from x towards the other vertex of e_i . Then $v_1, \dots, v_n$ are the vertices of a spherical polygon (called the link of P at x) which is convex since P is convex.

For each $i \in \{1, \dots, n\}$, the first-order deformation v of P determines a first-order deformation v' of the spherical polygon $(v_1, \dots, v_n)$, given at each vertex v_i by the vector $\dot{v}_i = dv(v_i)$; note that $dv(v_i)$ is orthogonal to v_i because v is an isometric deformation. Since v is isometric, it does not change (at first order) the interior angles of the faces of P ; since the interior angles of the faces of P at x are the lengths of the edges of $(v_1, \dots, v_n)$, the first-order deformation v' is isometric.

The angles $\alpha_1, \dots, \alpha_n$ of the polygon $(v_1, \dots, v_n)$ are equal to the dihedral angles $\theta(e_1), \dots, \theta(e_n)$ of P at the edges $e_1, \dots, e_n$. Therefore:

$$\begin{aligned} \sum_{i=1}^n W_{e_i} &= \sum_{i=1}^n (\langle dv(v_i), p_0 x \rangle + \langle v(x), v_i \rangle) \dot{\theta}_i \\ &= \sum_{i=1}^n (\langle \dot{v}_i, p_0 x \rangle + \langle v(x), v_i \rangle) \dot{\alpha}_i \\ &= \left\langle \sum_{i=1}^n \dot{\alpha}_i \dot{v}_i, p_0 x \right\rangle + \left\langle v(x), \sum_{i=1}^n \dot{\alpha}_i v_i \right\rangle . \end{aligned}$$

Since p_0 is in P and P is convex, $-p_0 x$ is in the positive cone over the interior of $(v_1, \dots, v_n)$, so that the first sum is non-positive by Theorem B_S , and it vanishes if and only if the deformation v' is trivial. The second sum is zero by Theorem A_S . So, by (1), the first sum is 0 for each vertex x of P , and none of the angles of P varies in the first-order deformation v , so that v is a trivial deformation.

Further results ? The proof of Theorem C given here has some flexibility. For instance, it should be possible to consider polyhedral surfaces with some singularities, e.g. ramifications points inside some of the faces. One can ask whether it also applies for polyhedral surfaces having a kind of ramification at the vertices, or to polyhedral surfaces which are not convex but satisfy a kind of weak convexity like the one considered in [Sto68, Kan90, RR00].

Another possible extension would be to convex surfaces with boundary, as considered e.g. in [Kan90]; in this setting it is necessary to have some constraints on the deformations at the boundary vertices. However, only the local convexity at the vertices plays a role, so that it should be possible to consider surfaces of higher Euler characteristic.

6 Convex polyhedral surfaces in the Minkowski space

This section contains an extension of Theorem C to a rigidity question for convex, equivariant surfaces in the Minkowski space, for which the techniques used here are particularly well adapted since there is a distinguished point. We first define the polyhedral surfaces that are considered. Then we basically follow the path taken above in the Euclidean case, adapting the proof from the sphere to the de Sitter plane. Along the way we recall some elementary facts of de Sitter geometry, proved here for completeness.

Equivariant embeddings of surfaces. Let S be a closed surface, which will be of genus at least 2 in all this section. We first define equivariant polyhedral embeddings of S . More general definitions could of course be given, but we stick to what is really necessary in our context.

Definition 6.1. An equivariant space-like polyhedral embedding of S in $\mathbb{R}_1^3$ is a couple (ϕ, ρ) , where:

- ϕ is a space-like polyhedral embedding of the universal cover $\tilde{S}$ of S in $\mathbb{R}_1^3$, i.e.:
 - ϕ is continuous.
 - There exists a cellular decomposition of S as the union of a finite number of cells, each the image by a diffeomorphism of the interior of a convex polygon in $\mathbb{R}^2$, such that the image by ϕ of each cell of the corresponding cellular decomposition of $\tilde{S}$ is the interior of a convex polygon in some space-like plane in $\mathbb{R}_1^3$.
 - For each space-like plane H in $\mathbb{R}_1^3$, the orthogonal projection of $\phi(\tilde{S})$ on H is one-to-one.
- ρ is a morphism from $\pi_1(S)$ to the isometry group of $\mathbb{R}_1^3$.
- For each $x \in \tilde{S}$ and each $\gamma \in \pi_1(S)$, $\phi(\gamma x) = \rho(\gamma)\phi(x)$.

We will say that (ϕ, ρ) is Fuchsian if the image of ρ is contained in the identity component of the sub-group of isometries fixing the origin, i.e. $\rho(\pi_1(S)) \subset SO_0(2, 1)$.

A direct consequence of the definition of an equivariant embedding of S in $\mathbb{R}_1^3$ is that the metric induced on $\tilde{S}$ by ϕ is invariant under the action of $\pi_1 S$, so that an equivariant embedding of S induces a metric on S .

Consider an equivariant embedding (ϕ, ρ) of S in $\mathbb{R}_1^3$. There is a natural notion of first-order deformation of (ϕ, ρ) ; it corresponds to the deformations of ϕ among the equivariant embeddings of S . Restricting our attention to the deformations among Fuchsian equivariant embeddings, such a deformation can be described as a couple $(\dot{\phi}, \dot{\rho})$, where:

- $\dot{\phi}$ is a vector field defined over $\phi(\tilde{S})$.
- $\dot{\rho} : \pi_1 S \rightarrow TSO(2, 1)$ is a map, such that:

$$\forall \gamma \in \pi_1 S, \dot{\rho}(\gamma) \in T_{\rho(\gamma)} SO_0(2, 1) ,$$

and, for all $\gamma, \gamma' \in \pi_1 S$, if we consider $\rho(\gamma)$ and $\rho(\gamma')$ as acting on $SO(2, 1)$ and $\dot{\rho}$ as sending elements of $\pi_1 S$ to vector fields on $SO(2, 1)$, we have:

$$\dot{\rho}(\gamma\gamma') = \dot{\rho}(\gamma)(\rho(\gamma')) + \rho(\gamma)_* \dot{\rho}(\gamma') .$$

- If we identify the elements of $so(2, 1)$ with the Killing fields on $\mathbb{R}_1^3$ which vanish at 0, then:

$$\forall x \in \tilde{S}, \forall \gamma \in \pi_1 S, \dot{\phi}(\gamma x) = \rho(\gamma)_* \dot{\phi}(x) + \dot{\rho}(\gamma)\phi(x) .$$

Fuchsian polyhedral surfaces. The main result of this section is the following infinitesimal rigidity statement. There is an analogous statement for smooth surfaces in [LS00].

Theorem 6.2. Let (ϕ, ρ) be a Fuchsian, convex equivariant embedding of a surface S (of genus at least 2) in $\mathbb{R}_1^3$. Let $(\dot{\phi}, \dot{\rho})$ be a first-order deformation of (ϕ, ρ) among Fuchsian equivariant embeddings. If the first-order variation of the metric induced on S by (ϕ, ρ) vanishes, then $(\dot{\phi}, \dot{\rho})$ is trivial, i.e. $\dot{\phi}$ is the restriction to $\phi(\tilde{S})$ of a Killing field of $\mathbb{R}_1^3$ which vanishes at 0, and $\dot{\rho} = 0$.

The proof comes after some preliminaries, basically following the proof of Theorem C and checking that the various parts of the proof carry over from the Euclidean to the Minkowski 3-dimensional space, and from the sphere to the de Sitter plane.

As in the Euclidean case, the main motivation for this theorem is that it should be a key point in the proof a statement describing the metrics induced on fuchsian equivariant polyhedra in $\mathbb{R}_1^3$, as in the Aleksandrov theorem quoted above for the Euclidean space. Namely, one would like to answer the following question:

Let S be a surface of genus at least 2, and let g be a flat metric on S with conical singularities, with total angle larger than 2π at each singular point. Is there a unique (up to global isometries) convex Fuchsian polyhedral embedding of S in $\mathbb{R}_1^3$ such that the induced metric is g ?

We do not answer this question here since it would demand some considerations leading us too far from Theorems A_S and B_S , the main theme of this text.

Distances and angles in the de Sitter plane. As mentioned in the introduction, the de Sitter plane, denoted by S_1^2 , is the quadric of equation $\langle x, x \rangle = 1$ in $\mathbb{R}_1^3$, with the induced metric. Details can be found e.g. in [Cox43, Cox57, O’N83].

Let x, x' be two points in S_1^2 . By analogy with the distance in the sphere $S^2 \subset \mathbb{R}^3$, we define the “distance” between x and x' as the number $d(x, y) \in \mathbb{C}/2\pi\mathbb{Z}$ such that:

$$\langle x, x' \rangle = \cos(d(x, x')) .$$

This leaves an indetermination concerning the sign of $d(x, x')$, which is removed by the following explicit description:

- $d(x, x') \in (0, \pi)$ when x and x' are distinct, non-antipodal points on a space-like geodesic.
- $d(x, x') = 0$ when x and x' are on a light-like geodesic.
- $d(x, x') \in i\mathbb{R}_+$ when x and x' are on a time-like geodesic.
- $d(x, x') \in \pi - i\mathbb{R}_+$ when x and x' are in two different connected components of the intersection of S_1^2 with a time-like plane containing 0 (each connected component of the intersection is then a time-like geodesic).

This definition differs from the one used in [Sch98] by a factor i , basically because here we stick as close as possible to spherical geometry, while the emphasis in [Sch98] was on hyperbolic geometry. The definition of [Sch98] would be obtained if we had taken the cosh of $d(x, x')$ instead of the cosine in the definition above. The proof below would then have less factors “ i ”, but it would be based on the hyperbolic rather than the spherical trigonometric formulas, thus blurring the analogy with the proof of Theorem C.

There is a related notion of angles in S_1^2 . Let $x \in S_1^2$, and let $u, v \in T_x S_1^2$ be such that $\langle u, u \rangle \neq 0$ and $\langle v, v \rangle \neq 0$; the angle between them is a number $\theta \in \mathbb{C}/2\pi\mathbb{Z}$ such that:

$$\langle u, v \rangle^2 = \cos^2(\theta) \langle u, u \rangle \langle v, v \rangle .$$

Note that θ is defined up to sign and up to the transformation $\theta \mapsto \pi - \theta$, which is quite normal since it depends on the direction of u and v and on whether we consider u or v first. This ambiguity which is removed by the following explicit description, in which we suppose that (u, v) is a positively oriented basis of $T_x S_1^2$:

- $\theta \in i\mathbb{R}_+$ if u and v are both future-oriented time-like vector (or if they are both past-oriented time-like vectors).
- $\theta \in \pi - i\mathbb{R}_+$ if u and v are time-like vectors, but u is future-oriented and v is past-oriented.
- $\theta \in i\mathbb{R}_+$ if u and v are both space-like vectors, and are in the same connected component of the set of space-like vectors at x .
- $\theta \in \pi - i\mathbb{R}_+$ if u and v are space-like vectors in different connected components of the set of space-like vectors at x .
- $\theta \in \pi/2 + i\mathbb{R}$ if u is space-like and v is time-like, or conversely.

Triangle formulas in the de Sitter plane. With those definitions, equation (1) and the sine formula hold in S_1^2 exactly as in the sphere.

Proposition 6.3. *Let (A, B, C) be a triangle in the de Sitter plane, with edge lengths a, b, c and angles α, β, γ . Suppose that $a, b, c \neq 0$. Then:*

$$\cos(a) = \cos(b) \cos(c) + \cos(\alpha) \sin(b) \sin(c) .$$

Moreover, if $a, b, c \neq 0$, then:

$$\frac{\sin(\alpha)}{\sin(a)} = \frac{\sin(\beta)}{\sin(b)} = \frac{\sin(\gamma)}{\sin(c)} .$$

Proof. Recall the definition of the cross-product $X \boxtimes Y$, in coordinates, used in the Minkowski space:

$$(x_1, x_2, x_3) \boxtimes (y_1, y_2, y_3) = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, -x_1 y_2 + x_2 y_1) .$$

Note that this definition is natural insofar as it can be obtained like the vector product in Euclidean space, by associating 1-forms to vectors, taking the wedge product, and associating a vector to the resulting 2-form. This shows that the definition given using coordinates is independent of the orthonormal basis of $\mathbb{R}_1^3$ which has been used.

It follows from this definition that, given 3 vectors X, Y and Z in $\mathbb{R}^3$, the number $\langle X, Y \boxtimes Z \rangle$ is the same whether the cross product and the scalar product are considered for the Euclidean or the Minkowski space structure (there are two sign differences but they cancel out). Therefore, it remains true in the Minkowski space that:

$$\langle X, Y \boxtimes Z \rangle = \langle Y, Z \boxtimes X \rangle = \langle Z, X \boxtimes Y \rangle .$$

The same argument also shows that $\langle X, X \boxtimes Y \rangle = \langle Y, X \boxtimes Y \rangle = 0$ in the Minkowski space, so that $X \boxtimes Y$ is orthogonal (for the Minkowski scalar product) to X and to Y .

It is also easy to check, by taking a “good” orthonormal basis of $\mathbb{R}_1^3$, that, if X is a space-like unit vector orthogonal to Y and Z , then:

$$\langle X \boxtimes Y, X \boxtimes Z \rangle = \langle Y, Z \rangle .$$

Moreover, let $X, Y \in S_1^2$ be two non-collinear vectors of $\mathbb{R}_1^3$, there exists a direct orthonormal basis (X, Y', Z) of $\mathbb{R}_1^3$, such that $Y = \cos(d(X, Y))X + \sin(d(X, Y))Y'$; a direct computation then shows that:

$$X \boxtimes Y = \sin(d(X, Y))Z .$$

Now let $A, B, C \in S_1^2$ be the 3 vertices of a de Sitter triangle. Then B and C can be decomposed as $B = B_{\parallel}A + B_{\perp}$ and $C = C_{\parallel}A + C_{\perp}$, where $B_{\parallel}, C_{\parallel} \in \mathbb{R}$ and $B_{\perp}, C_{\perp} \in \mathbb{R}_1^3$ are vectors orthogonal to A . Then:

$$\begin{aligned} \langle A \boxtimes B, A \boxtimes C \rangle &= \langle A \boxtimes (B_{\parallel}A + B_{\perp}), A \boxtimes (C_{\parallel}A + C_{\perp}) \rangle = \langle A \boxtimes B_{\perp}, A \boxtimes C_{\perp} \rangle = \\ &= \langle B_{\perp}, C_{\perp} \rangle = \langle B, C \rangle - B_{\parallel}C_{\parallel} \langle A, A \rangle = \langle B, C \rangle - \langle A, B \rangle \langle A, C \rangle . \end{aligned}$$

But $A \boxtimes B = \sin(c)N_C$ and $A \boxtimes C = \sin(b)N_B$, where N_C and N_B are the unit vectors orthogonal to the (oriented) plane containing $0, A, B$ and $0, A, C$, respectively. By definition, $\langle N_C, N_B \rangle = \cos(\alpha)$, and we find that:

$$\cos(\alpha) \sin(b) \sin(c) = \langle A \boxtimes B, A \boxtimes C \rangle = \langle B, C \rangle - \langle A, B \rangle \langle A, C \rangle = \cos(a) - \cos(b) \cos(c) ,$$

which proves the first equation.

To prove the sine formula, note that N_B and N_C are both orthogonal to A , so that $(A \boxtimes B) \boxtimes (A \boxtimes C)$ is collinear to A . It follows that:

$$\langle (A \boxtimes B) \boxtimes (A \boxtimes C), A \rangle = \sin(c) \sin(b) \sin(\alpha) .$$

But, using the same decomposition of B and C as above, we have:

$$(A \boxtimes B) \boxtimes (A \boxtimes C) = (A \boxtimes B_{\perp}) \boxtimes (A \boxtimes C_{\perp}) = B_{\perp} \boxtimes C_{\perp} = B \boxtimes C - B_{\parallel}(A \boxtimes C_{\perp}) - C_{\parallel}(A \boxtimes B_{\perp}) ;$$

taking the scalar product with A yields:

$$\langle (A \boxtimes B) \boxtimes (A \boxtimes C), A \rangle = \langle B \boxtimes C, A \rangle .$$

It follows that the quantity $\sin(b) \sin(c) \sin(\alpha)$ is invariant under a cyclic permutation on (a, b, c) and (α, β, γ) , and the second equation in the proposition follows. $\square$

Polygons in the de Sitter plane. The analog of Theorem A_S and Theorem A_H also holds for de Sitter polygons with non-degenerate edges (when one of the edges is degenerate, the notion of angle is not well-defined, so that the statement would not make sense).

Theorem A_{dS} . *Let $p = (v_1, \dots, v_n)$ be a de Sitter polygon with non-degenerate edges. Let $\alpha_1, \dots, \alpha_n$ be its angles, and let $\dot{\alpha}_1, \dots, \dot{\alpha}_n \in \mathbb{R}$ be a first-order variation of its angles induced by an isometric first-order deformation of p . Then:*

$$\sum_{i=1}^n \dot{\alpha}_i v_i = 0 ,$$

where the v_i are considered as points in $S_1^2 \subset \mathbb{R}_1^3$.

Conversely, if this equation is satisfied by an n -uple $(\dot{\alpha}_1, \dots, \dot{\alpha}_n)$ and moreover the v_i are not all on a de Sitter geodesic – or on the intersection of S_1^2 with a Minkowski plane containing the origin – then there exists an isometric first-order deformation of p such that the $\dot{\alpha}_i$ are the associated first-order variations of the α_i .

The proof follows exactly the proof of Theorem A_H .

Again as in the sphere, Theorem A_{dS} leads to the definition of a quadratic invariant b defined on the first-order isometric deformations of a de Sitter polygon (with non-degenerate edges). b is defined as:

$$b(U) = \sum_{i=1}^n d\alpha_i(U) dv_i(U) ,$$

As in the spherical and the hyperbolic case, if two first-order infinitesimal deformations U and U' differ by a trivial deformation, then $b(U) = b(U')$. However, the definition of the angles which we have used means that $b(U)$ is imaginary.

The positivity of b . In the same way, the “positivity” property of b which we found in the spherical case still holds for the de Sitter polygons which are duals of convex hyperbolic polygons.

Theorem B_{dS} . *Let p be a convex polygon in S_1^2 , which is dual to a convex hyperbolic polygon p^* , and let U be a non-trivial infinitesimal first-order deformation of p . Then $ib(U) \in \mathbb{R}_+^* \text{int}(p^*)$, i.e. $ib(U)$ is contained in the positive cone over the interior of p^* .*

The proof follows the proof of Theorem B_S , but some additional details are necessary. Using Proposition 6.3, the computations done in section 3 carry over to the de Sitter case. In particular, the scalar product of $b(U)$ with v_1 remains diagonal in the basis $(U_2, \dots, U_{n-2})$. Equation (1) can be rewritten in a way which is more convenient for us, as:

$$\langle b(U_i), v_1 \rangle = \frac{\sin^2(d_{1,i+1}) \sin(\alpha'_1)}{\sin(d_{0,i+1}) \sin(d_{i,i+1}) \sin(\alpha'_0) \sin(\alpha'_i)} (dd_{1,i+1}(U_i))^2 .$$

This follows from (1) because, in Lemma 3.3, the definition of V is that, under this first-order variation, the distance $d_{1,i+1}$ between v_1 and v_{i+1} varies at speed 1; this also uses the fact that b is a quadratic form. Recall that, in this equation α'_0, α'_1 and α'_i are not the angles of the polygon p , but rather the angles of the quadrilateral (v_0, v_1, v_i, v_{i+1}) at the vertices v_0, v_1 and v_i .

Using as variable the cos of the distance from v_1 and v_{i+1} , we obtain:

$$\langle b(U_i), v_1 \rangle = \frac{\sin(\alpha'_1)}{\sin(d_{0,i+1}) \sin(d_{i,i+1}) \sin(\alpha'_0) \sin(\alpha'_i)} (d\langle v_1, v_{i+1} \rangle(U_i))^2 .$$

Clearly, $(d\langle v_1, v_{i+1} \rangle(U_i))^2 \in \mathbb{R}_+$; moreover, since the edges of p are space-like, $d_{i,i+1} \in (0, \pi)$, so that $\sin(d_{i,i+1}) \in (0, 1)$. However, to understand the signs of the various sines appearing in this equation, it is necessary to consider four different cases, depending on whether $d_{1,i}$ and $d_{0,i+1}$ are in $(0, \pi)$ or in $\pi - i\mathbb{R}_+^*$, and then the limit cases. The four cases are shown, in the projective model of half of S_1^2 , in Figure 2.

1st case: $d_{1,i} \in (0, \pi), d_{0,i+1} \in (0, \pi)$. In other terms, v_0 and v_{i+1} are on a space-like geodesic, and v_1 and v_i are on a space-like geodesic. Then $\sin(d_{0,i+1}) \in (0, 1)$. Moreover, the angles α'_0, α'_1 and α'_i are all




Figure 3: Four possible cases.

of the form $\pi - ir$, for some $r > 0$, so that $\sin(\alpha'_0), \sin(\alpha'_1)$ and $\sin(\alpha'_i)$ are in $i\mathbb{R}_+^*$. So equation (1) shows that $\langle b(U_i), v_1 \rangle \in -i\mathbb{R}_+^*$.

2nd case: $d_{1,i} \in (0, \pi), d_{0,i+1} \in \pi - i\mathbb{R}_+^*$. This means that v_1 and v_i are on a space-like geodesic, while v_0 and v_{i+1} are on the intersection with S_1^2 of a time-like plane in $\mathbb{R}_1^3$ containing 0 (but on different connected components of this intersection). Now $\sin(d_{0,i+1}) \in i\mathbb{R}_+^*$. α'_1 and α'_i are of the form $\pi - ir$ for some $r > 0$, so that $\sin(\alpha'_1)$ and $\sin(\alpha'_i)$ are in $i\mathbb{R}_+^*$, but α'_0 is of the form $\pi/2 + ir$, for some $r \in \mathbb{R}$, and it follows that $\sin(\alpha'_i) = \cos(ir) = \cosh(r) \in [1, \infty)$. Therefore, (1) again shows that $\langle b(U_i), v_1 \rangle \in -i\mathbb{R}_+^*$.

3rd case: $d_{1,i} \in \pi - i\mathbb{R}_+^*, d_{0,i+1} \in (0, \pi)$. Then $\sin(d_{0,i+1}) \in (0, 1)$. Moreover, α'_0 is of the form $\pi - ir$, for some $r > 0$, while α'_1 and α'_i are of the form $\pi/2 + ir$, for some $r \in \mathbb{R}$. So $\sin(\alpha'_0) \in i\mathbb{R}_+^*$, while $\sin(\alpha'_1), \sin(\alpha'_i) \in [1, \infty)$. By (1), $\langle b(U_i), v_1 \rangle \in -i\mathbb{R}_+^*$ also in this case.

4th case: $d_{1,i} \in \pi - i\mathbb{R}_+^*, d_{0,i+1} \in \pi - i\mathbb{R}_+^*$. In this case, $\sin(d_{0,i+1}) \in i\mathbb{R}_+^*$, and α'_0, α'_1 and α'_i are of the form $\pi/2 + ir$, for some $r \in \mathbb{R}$. Therefore, $\sin(\alpha'_0), \sin(\alpha'_1)$ and $\sin(\alpha'_i)$ are in $[1, \infty)$, and, again, $\langle b(U_i), v_1 \rangle \in -i\mathbb{R}_+^*$.

Last case: v_1 and v_i are in the intersection with S_1^2 of a light-like plane in S_1^2 containing 0, or the same holds of v_0 and v_{i+1} . The result follows from an approximation of the quadrilateral (v_0, v_1, v_i, v_{i+1}) by a sequence of quadrilaterals which are in one of the four cases detailed above, and a corresponding sequence of approximations of U_i .

Summing over $i \in \{2, \dots, n-2\}$, we find that:

$$\langle ib(U), v_1 \rangle = \sum_{i=2}^{n-2} i \langle b(U_i), v_1 \rangle > 0.$$

The same holds with v_1 replaced by any of the other vertices of p , and Theorem B_{dS} follows.

Proof of the rigidity theorem. The proof of Theorem 6.2 follows quite precisely the proof of Theorem C. We consider a Fuchsian, equivariant, convex polyhedral embedding (ϕ, ρ) of a surface S of genus at least 2 in $\mathbb{R}_1^3$. Let $p_0 = 0 \in \mathbb{R}_1^3$, i.e. p_0 is the point which is fixed by ρ . We call u_0 the function, defined on $\mathbb{R}_1^3$, as: $u_0(x) = \langle x, x \rangle / 2$. Then, since 0 is fixed by the representation ρ , the restriction of u_0 to $\phi(\tilde{S})$ is invariant under the action of $\pi_1 S$ by ρ , so that u_0 defines a function over S .

Let $(\dot{\phi}, \dot{\rho})$ be a first-order deformation of (ϕ, ρ) among Fuchsian embeddings. Let e be an oriented edge of $\phi(\tilde{S})$, let θ_e be the corresponding dihedral angle, and let $\dot{\theta}_e$ be the first-order variation of this dihedral angle under the first-order variation $(\dot{\phi}, \dot{\rho})$. We associate to e the number W_e defined as: $W_e = \dot{\theta}_e du_0(p'(t))$, where $p(t)$ is a parametrization of e at speed one. As in the Euclidean case – and for the same reasons – W_e is independent of t . By construction, this quantity is invariant under the action of $\pi_1 S$.

Again in this setting, $\sum_v \sum_{e_- = v} W_e = 0$. On the other hand, let x be a vertex of $\phi(\tilde{S})$. Let $e_1, \dots, e_n$ be the oriented edges starting from x , in the cyclic order in which they appear. The link of $\phi(\tilde{S})$ at x is a convex space-like polygon p , with vertices $v_1, \dots, v_n$, corresponding to $e_1, \dots, e_n$, which is the dual of a hyperbolic polygon. The first-order isometric deformation $(\dot{\phi}, \dot{\rho})$ induces a first-order deformation U of p . Since $(\dot{\phi}, \dot{\rho})$ is isometric, it does not change (at first order) the interior angles of the faces of $\phi(\tilde{S})$ adjacent to x , so that the first-order deformation U is isometric.

As in the Euclidean case, if $\dot{v}_i$ is the first-order displacement of v_i under U and if α_i is the angle of p at v_i , then:

$$\begin{aligned} \sum_{i=1}^n W_{e_i} &= \sum_{i=1}^n \left(\langle d\dot{\phi}(v_i), x \rangle + \langle \dot{\phi}(x), v_i \rangle \right) \dot{\theta}_{e_i} \\ &= \sum_{i=1}^n \left(\langle \dot{v}_i, x \rangle + \langle \dot{\phi}(x), v_i \rangle \right) \dot{\alpha}_i \\ &= \left\langle \sum_{i=1}^n \dot{\alpha}_i \dot{v}_i, x \right\rangle + \left\langle \dot{\phi}(x), \sum_{i=1}^n \dot{\alpha}_i v_i \right\rangle. \end{aligned}$$

The second terme vanishes by Theorem A_{ds}. By convexity of $\phi(\tilde{S})$, x is in the interior of p , while, by Theorem B_{ds}, $ib(U)$ is in the interior of the dual polygon p^* . It follows that $i \sum_{i=1}^n W_{e_i} \geq 0$, with equality if and only if the first-order deformation U is trivial. The end of the proof is the same as in the Euclidean case.

Note that the rigidity argument given here for equivariant polyhedral embeddings in $\mathbb{R}_1^3$ could perhaps be used in other situations, for instance for closed, convex polyhedra in $\mathbb{R}_1^3$ which have faces which are not necessarily space-like. However the infinitesimal rigidity of convex polyhedra in $\mathbb{R}_1^3$ – even when some faces are not space-like – follows from the infinitesimal rigidity of convex polyhedra in the Euclidean space (see e.g. [Sch01]) – so that having a direct Minkowski proof is not so important.

7 Natural metrics on the moduli space of polygons

The spaces of polygons in the sphere (or in other constant curvature spaces) are of interest in topological or algebraic terms (see e.g. [KM95, KM99]) but also in metric terms. This is in particular true of spaces of convex Euclidean polygons with fixed angles (rather than fixed edge lengths); Bavard and Ghys [BG92] showed that those spaces can be naturally identified with hyperbolic polyhedra, which are Coxeter polyhedra when some fairly simple conditions on the angles of the polygons are satisfied. This is related to the construction by Thurston [Thu98] of complex hyperbolic orbifolds as moduli spaces of flat metrics with conical singularities on the sphere (see [Fil]), and also related to [DM86].

The positivity property of b stated in Theorem B_S can be used to define some natural metrics on the spaces of convex polygons in the sphere (or in the hyperbolic plane) with given edge lengths. We will describe those constructions here, and indicate how they appear to be related to the metric used by Bavard and Ghys on spaces of Euclidean polygons with given angles, which is recovered as a limit case. The finer properties of the metrics on the spaces of spherical or hyperbolic polygons are not studied here.

Definitions of some metrics. The definitions of the metrics we want to consider stem from the following elementary remark. We call $P_{n,S}^c$ the space of convex polygons with n vertices in S^2 , and $\mathcal{P}_{n,S}^c$ the quotient of $P_{n,S}^c$ by $SO(3)$. Given an n -uple $(l_1, \dots, l_n) \in \mathbb{R}_+$, we call $P_S^c(l)$ the space of convex polygons in S^2 with edge lengths equal to l , and $\mathcal{P}_S^c(l) := P_S^c(l)/SO(3)$.

Remark 7.1. Let $F : P_{n,S}^c \rightarrow S^2$ be a map such that:

- For all $p \in P_{n,S}^c$, $F(p)$ is in the interior of p .
- For all $p \in P_{n,S}^c$ and $\gamma \in SO(3)$, $F(\gamma p) = \gamma F(p)$.

Let $p_0 \in P_{n,S}^c$ and let l be its edge lengths. Then the bilinear form $\langle b_2(\cdot, \cdot), F(p) \rangle$ on the tangent space to $P_S^c(l)$ at each point p defines a Riemannian metric on $P_S^c(l)$, which is compatible with the quotient by $SO(3)$. Therefore it defines a Riemannian metric on $\mathcal{P}_S^c(l)$.

Proof. By Theorem B_S , for each $p \in P_S^c(l)$ and each non-trivial deformation $U \in T_p P_S^c(l)$, $b(U) \in \mathbb{R}_+^* \text{int}(p^*)$. This means precisely that the scalar product of $b(U)$ with any point in the interior of p is positive, and this holds in particular of the scalar product of $b(U)$ with $F(p)$, so that $\langle b_2(\cdot, \cdot), F(p) \rangle$ defines a positive semi-definite bilinear form on $P_S^c(l)$, and the kernel corresponds precisely to the trivial deformations.

We have already seen that, if U' is a trivial first-order deformation of p , then $b(U + U') = b(U)$. The trivial first-order deformations of p , seen as vectors in $T_p P_S^c(l)$, are the elements of the kernel of the differential of the projection from $T_p P_S^c(l)$ to $T_{[p]} \mathcal{P}_S^c(l)$, where $[p]$ is the image of p under the quotient of $P_S^c(l)$ by $SO(3)$. So the Riemannian metric defined by $\langle b_2(\cdot, \cdot), F(p) \rangle$ on $T_p P_S^c(l)$ is the pull-back by the projection of a Riemannian metric on $T_{[p]} \mathcal{P}_S^c(l)$.

Now let $\gamma \in SO(3)$, and let U be a first-order deformation of p . Considering γ as a map acting on $P_S^c(l)$, we associate to U a first-order deformation $(d_p \gamma)(U)$ of γp . The definition of b shows that it is “invariant” under the action of γ , i.e. that $b((d_p \gamma)(U)) = \gamma b(U)$. It follows that:

$$\langle b((d_p \gamma)(U)), F(\gamma p) \rangle = \langle \gamma b(U), \gamma F(p) \rangle = \langle b(U), F(p) \rangle ,$$

so that the bilinear form $\langle b_2(\cdot, \cdot), F(p) \rangle$ is invariant under the action of $SO(3)$. $\square$

There are several possible choices for the function F . It is quite natural to take some kind of barycenter of p , for instance:

- The barycenter of the vertices of p (all with the same weight), which we will call $C_v(p)$. This is the most obvious solution, however we will see below that it lacks one desirable property.
- The barycenter of the area form on the interior of p , which we call $C_i(p)$. This is another quite obvious choice, we will see below that it has some better properties.
- The barycenter of the vertices of p , with weights equal to the exterior angles, which we call $C_\alpha(p)$.
- The barycenter of p itself, i.e. of the union of its edges, denoted here by $C_\partial(P)$.

Given a family of lengths $l = (l_1, \dots, l_n)$, we will call $g_v(l)$ (resp. $g_i(l)$, $g_\alpha(l)$, $g_\partial(l)$) the Riemannian metrics on $\mathcal{P}_S^c(l)$ described by Remark 7.1 with F equal to C_v (resp. to C_i , C_α , C_∂).

A geometric property of the barycenter of the area. The drawback with the choice of the barycenter of the vertices of p is that it does not have the following property, which is satisfied by the barycenter of the interior of p .

Proposition 7.2. Let p be a convex polygon in S^2 . The barycenter $C_i(p)$ of the interior of p is contained in the interior of the dual polygon p^* .

Proof. By definition, $C_i(p)$ is in the interior of p^* if and only if, for all point x in the interior of p , $\langle C_i(p), x \rangle \geq 0$. This is equivalent to the fact that $C_i(p)$ is at distance at most $\pi/2$ from any point in p . Since the space of convex polygons with n vertices is connected, the result follows from the following assertion: if p is contained in the hemisphere centered at $C_i(p)$ and one point of p is in the boundary of this hemisphere, then p is a polygon with 2 edges, which are both one half of a great circle, with one endpoint on x .

So we suppose that p is contained in the hemisphere H centered at $C_i(p)$, with one point x in the boundary, and that it has more than two edges. Then the connected component of x in the intersection of p with ∂H is either only one point x , or it an edge of p . We first consider the first case.

Consider the projective model of the hemisphere H , obtained by projecting the points of H radially to the plane in $\mathbb{R}^3$ tangent to S^2 at $C_i(p)$. The image of p in this model is a “non-compact” convex polygon which we call p_e , i.e. the boundary of a non-compact polygonal domain, with two parallel infinite edges e and e' , which is the image of the interior of p .

The projective model has the property that the symmetry in H with respect to the geodesic line g at distance $\pi/2$ from x (which contains $C_i(p)$) acts like the symmetry in $\mathbb{R}^2$ with respect to the line g_e containing the origin and orthogonal to e and to e' . But a simple convexity argument shows that, if $y \in \mathbb{R}^2$ is on the same side of g_e as the infinite ends of e and e' , and if y is not in the interior of p_e , then the image y' of y under the symmetry with respect to g_e is not in the interior of p_e ; otherwise, some support line of p_e would go between y and y' , and then it would have to “cut” the infinite edges e and e' . Moreover, p_e is not invariant under the symmetry with respect to g_e , unless it is an infinite strip, and then p is a polygon with two edges, each of which is half of a great circle.

It follows that the same statement is true in H : if a point z is on the same side of g as x , but is not contained in the interior of p , then the image of z under the symmetry with respect to g is not contained in the interior of p . Thus:

$$\int_{\text{int}(p)} \langle x, \cdot \rangle da > 0 ,$$

and this contradicts the fact that $C_i(p)$, which is orthogonal to x , is the barycenter of the interior of p . This shows the result when x is a vertex of p .

If the intersection of p with ∂H contains an edge of p , the same argument can be used, taking as x one point of this edge. The only difference is that the two infinite edges of p_e are not parallel; the key point remains that, if H_+ is the hemisphere bounded by g and containing x and H_- is the other hemisphere bounded by g , then $H_- \cap \text{int}(p)$ is contained in the image under the symmetry with respect to g of $H_+ \cap \text{int}(p)$. The proof can again be obtained using the projective model of H . We leave the details to the reader. $\square$

Note that this statement is not correct with $C_i(p)$ replaced with $C_v(p)$, however it might hold with $C_i(p)$ replaced by $C_\alpha(p)$ or $C_\partial(p)$.

Euclidean polygons in the limit. Let $\alpha = (\alpha_1, \dots, \alpha_n)$ be a set of positive numbers with sum equal to 2π . We call $\mathcal{P}_E^*(\alpha)$ the space of convex Euclidean polygons with exterior angles equal to $\alpha_1, \dots, \alpha_n$, considered up to the Euclidean isometries. We denote by $\mathcal{P}_E^{*,A}(\alpha)$ the subspace of $\mathcal{P}_E^*(\alpha)$ of polygons of area equal to A .

There is a natural metric on $\mathcal{P}_E^{*,1}(\alpha)$, defined by Bavard and Ghys [BG92], called the “area form”, which we will denote by g_A here. It can be defined by considering the convex polygons with edges parallel to some given directions, up to translation (this is very close to considering polygons with given angles, up to isometry), and noting that the area is a quadratic form over the space of those polygons. Bavard and Ghys show that the signature of this form is $(1, n-3)$, and the moduli space of Euclidean polygons with given angles (up to the homotheties), with the induced metric, is isometric to the interior of a finite volume hyperbolic polyhedron. Moreover, under some explicit conditions on α , this polyhedron is Coxeter, i.e. the group generated by the reflections in its faces is discrete. There is a relationship between those metrics on spaces of convex Euclidean polygons with given angles and the metrics defined above on spaces of spherical polygons with given edge lengths.

Theorem 7.3. *Let $(l^k)_{k \in \mathbb{N}} = (l_1^k, \dots, l_n^k)_{k \in \mathbb{N}}$ be a sequence of n -uples of positive numbers. Suppose that $\lim_{k \rightarrow \infty} l^k = \alpha$, with $\alpha = (\alpha_1, \dots, \alpha_n)$ such that $\sum_i \alpha_i = 2\pi$ and that there is no $p, q \in \{1, \dots, n\}$ such that $\sum_{i=p}^q \alpha_i = \pi$. Then (by [BG92]), $\mathcal{P}_E^c(\alpha)$ is compact, and:*

$$(\mathcal{P}_S^c(l^k), \frac{1}{2a_k} g_i(l^k)) \rightarrow (\mathcal{P}_E^{*,1}(\alpha), g_A) ,$$

where $a_k := 2\pi - \sum_{i=1}^n l_i^k$, and the convergence is Lipschitz.

The proof is given below, it uses some preliminary statements.

Remark 7.4. *Let $l = (l_1, \dots, l_n)$ be a family of positive numbers.*

1. Suppose that there exists a convex polygon in S^2 with edge lengths given by l , then the sum of the l_i is less than 2π .
2. If p is a convex spherical polygon with edge lengths given by l , then p^* has exterior angles given by l , and area equal to $2\pi - \sum l_i$.
3. Suppose that $(p_k)_{k \in \mathbb{N}}$ is a sequence of convex spherical polygons, with p_k of edge lengths $l^k = (l_1^k, \dots, l_n^k)$ with $\lim_{k \rightarrow \infty} l_i^k = l_i$ and $\sum_{i=1}^n l_i = 2\pi$. Then, after taking a subsequence, $(p_k)_{k \in \mathbb{N}}$, considered as a sequence of subsets of S^2 , converges either to a great circle, or to a polygon with two edges, each being one half of a great circle. This second case can happen only if there exist $p, q \in \{1, \dots, n\}$ such that $\sum_{i=p}^q l_i = \pi$.

Proof. The proof of the first two points is either elementary or well-known. The third point follows from the second, because the sequence of dual polygons has area going to 0, so that, after taking a subsequence, the dual polygons converge either to a point, or to a segment. In the first case, (p_k) converges to a great circle, while, in the second case, (p_k) converges to a polygon with two edges, both of which is one half of the great circle which is dual to one of the endpoints of the segment. If the convergence is to a segment, then the limits of the exterior angles of the dual polygon at the vertices converging to each vertex of the segment has to sum to π , and the condition on the lengths of the edges follows. $\square$

We now need an additional notation concerning spaces of polygons with fixed angles in the sphere. Given $\alpha = (\alpha_1, \dots, \alpha_n)$, we call $\mathcal{P}_S^*(\alpha)$ the space of convex polygons in S^2 with angles given by α , up to the global isometries. There is a natural map Δ from $\mathcal{P}_S^c(\alpha)$ to $\mathcal{P}_S^*(\alpha)$ (and conversely), defined by sending a convex polygon p to the dual polygon p^* . Therefore, we can consider on $\mathcal{P}_S^*(\alpha)$ the pull-back metric Δ^*g_i , which we will call g_i^* .

It is actually helpful to consider g_i and g_i^* as symmetric bilinear forms on the space of all convex polygons in S^2 with n vertices, $\mathcal{P}_{S,n}^c$. Then g_i is defined, as for the more restricted space of polygons with given edge lengths, as:

$$g_i(U, V) = \frac{1}{2} \sum_{i=1}^n \langle d\alpha_i(U) dv_i(V) + d\alpha_i(V) dv_i(U), C_i(p) \rangle ,$$

and $g_i^* := \Delta^*g_i$. In the same way, we define the area form, already mentioned above, as a symmetric bilinear form over the space of convex Euclidean polygons with n vertices. For any point x_0 in the interior of p , the area form can be defined as:

$$g_A(U, V) = \frac{1}{4} \sum_{i=1}^n dl_i(U) dh_i(V) + dl_i(V) dh_i(U) ,$$

where h_i is the distance from x_0 to the i^{th} edge.

Let $q = (v_1, \dots, v_n)$ be a convex polygon in S^2 . Let $u \in SO(3)$ be an isometry sending its barycenter $C_v(q)$ to the “north pole” $(0, 0, 1) \in S^2$, and let $\rho(q)$ be the image of $u(q)$ by the projective map from the upper hemisphere to $\mathbb{R}^2$. If q is such that $C_v(q)$ is in the interior of q^* , then $u(q)$ is contained in the upper hemisphere, so that $\rho(q)$ is a convex Euclidean polygon. It is easy to check that $\rho(p)$, considered up to the Euclidean isometries, is independent of the choice of u (which is uniquely defined up to a rotation). We call $\bar{\rho}$ the map sending q to $\rho(q)$. Moreover, given $q' \in \mathcal{P}_E^c$, we call $(\alpha_1^E, \dots, \alpha_n^E)$ its angles, so that the α_i^E , for $1 \leq i \leq n$, define n functions on $\mathcal{P}_E^c$.

Proposition 7.5. *Let $(\alpha_1, \dots, \alpha_n)$ be such that $\sum_i \alpha_i = 2\pi$, but such that there is no $p, q \in \{1, 2, \dots, n\}$ with $\sum_{i=p}^q \alpha_i = \pi$. Let $(\alpha^k)_{k \in \mathbb{N}}$ be a sequence of n -uples of positive numbers such that, for all $k \in \mathbb{N}$, $\sum_i \alpha_i^k < 2\pi$, and that $\lim_{k \rightarrow \infty} \alpha^k = \alpha$. Then there exists a sequence $(\epsilon_k)_{k \in \mathbb{N}} \rightarrow 0$ such that, for all $k \in \mathbb{N}$:*

1. over $\mathcal{P}_S^*(\alpha^k)$, $(1 - \epsilon_k)g_i^* \leq 2\bar{\rho}^*g_A \leq (1 + \epsilon_k)g_i^*$.
2. for all $1 \leq i \leq n$, $\|d(\alpha_{i|\bar{\rho}(\mathcal{P}_S^*(\alpha^k))}^E)\|_{g_A} \leq \epsilon_k$.

Proof. Let $q \in \mathcal{P}_{S,n}^c$, and let $e_1, \dots, e_n$ be the edges of q . Let $C_0 := C_i(q^*)$, we consider here C_0 as a fixed point in S^2 , even when considering first-order deformations of q^* . For each $i \in \{1, 2, \dots, n\}$, let β_i be the spherical distance from C_0 to e_i . Consider a first-order deformation U of q , it determines a first-order deformation U^* of the dual polygon q^* . If v_i^* is the vertex of q^* which is dual to e_i , we have:

$$\langle dv_i^*(U^*), C_i(q^*) \rangle = d(\sin(\beta_i))(U) = \cos(\beta_i) d\beta_i(U) .$$

Since the edge lengths of q are the exterior angles of q^* , it follows that, calling l_i the edge lengths of q , we have:

$$\langle b(U^*), C_i(q^*) \rangle = \sum_{i=1}^n \cos(\beta_i) dl_i(U) d\beta_i(U) ,$$

so that:

$$g_i^*(U, U) = \sum_{i=1}^n \cos(\beta_i) dl_i(U) d\beta_i(U) .$$

We now add to U a trivial deformation, so as to obtain that the barycenter $C_v(q)$ is fixed by U , and also that one of the vertices of q , for instance v_1 , is not moved. There exists a constant $C > 0$ (depending on n) such that, for all $1 \leq i \leq n$:

$$\|dv_i(U)\| \leq C \left(\sum_j |dl_j(U)| + |d\alpha_j(U)| \right) .$$

Let $q^E := \rho(q)$, let l_i^E be the edge lengths of q^E , and let h_i be the distances between the edges of q^E and $\rho(C_0)$. Let U^E be the first-order deformation of q^E which is induced by U through ρ . We now suppose that $q \in \mathcal{P}_S^*(\alpha^k)$ (for some $k \in \mathbb{N}$). It follows from Remark 7.4 that, if k is large enough, q is contained in the ball centered at $C_i(q^*)$ of radius λ_k , where $\lim_{k \rightarrow \infty} \lambda_k = 0$. But the projective map ρ is smooth, and its differential at the “north pole” is an isometry. Thus it follows from (1) that, for k large enough, we have, for some constant $C > 0$:

$$\begin{aligned} |dl_i^E(U^E) - dl_i(U)| &\leq C\lambda_k \left(\sum_j |dl_j(U)| + |d\alpha_j(U)| \right) , \\ |dh_i(U^E) - d\beta_i(U)| &\leq C\lambda_k \left(\sum_j |dl_j(U)| + |d\alpha_j(U)| \right) . \end{aligned}$$

Moreover, the same argument – using the fact that $d\rho$ is conformal at the “north pole” – yields that:

$$|d\alpha_i^E(U^E) - d\alpha_i(U)| \leq C\lambda_k \left(\sum_j |dl_j(U)| + |d\alpha_j(U)| \right) .$$

Using the description of g_i^* and the definition of g_A given above, we obtain with (1) and (1) that, for another constant $C' > 0$:

$$|g_A(U^E, U^E) - g_i^*(U, U)| \leq C'\lambda_k \left(\sum_j |dl_j(U)| + |d\alpha_j(U)| \right)^2 .$$

On the other hand, we have seen in section 3 that $\langle b, C_i(q^*) \rangle$ is positive definite on the tangent plane at each point of $\mathcal{P}_S^c(\alpha^k)$; it follows by compactness that there exists a constant $C'' > 0$ such that, for all $k \in \mathbb{N}$, all $q \in \mathcal{P}_S^*(\alpha^k)$ and all first-order deformation U of q leaving the angles constant at first order:

$$g_i^*(U, U) \geq \frac{1}{C''} \left(\sum_j |dl_j(U)| + |d\alpha_j(U)| \right)^2 .$$

The first point of the proposition follows from this equation and from (1), while the second point then follows from (1). $\square$

Proof of Theorem 7.3. The first point of Proposition 7.5 shows that the restriction of the $\bar{\rho}$ to $\mathcal{P}_S^*(\alpha^k)$ is a Lipschitz map, which multiplies the metric by 2 (up to a factor converging to 1). The polygons which are in the image are, by Remark 7.4, close to 0 in $\mathbb{R}^2$. Since ρ is an isometry at the “north pole”, it follows that their area is equivalent to a_k as $k \rightarrow \infty$.

Let H_k be the homothety of ratio $1/\sqrt{a_k}$, acting on Euclidean polygons. It follows from the above argument that $H_k \circ \bar{\rho}(\mathcal{P}_S^*(\alpha^k))$ converges to $\mathcal{P}_E^{*,1}(\alpha)$. Moreover, the second point of Proposition 7.5 shows that the norm of the differential of the functions α_i^E converges to 0 on the image, so that the $H_k \circ \bar{\rho}(\mathcal{P}_S^*(\alpha^k))$ actually converge C^1 to $\mathcal{P}_E^{*,1}(\alpha)$, and therefore the metric induced on the $H_k \circ \bar{\rho}(\mathcal{P}_S^*(\alpha^k))$ by g_A converge to the metric on $\mathcal{P}_E^{*,1}(\alpha)$.

But the first point of Proposition 7.5 shows that the metrics induced on the $H_k \circ \bar{\rho}(\mathcal{P}_S^*(\alpha^k))$ are close to $(1/2a_k)g_i^*$, and the proof of the theorem follows. $\square$

A statement analog to Theorem 7.3 appears to hold when there exist $p, q \in \{1, \dots, n\}$ such that $\sum_{i=p}^q \alpha_i = \pi$. In that case, however, $\mathcal{P}_E^{*,1}(\alpha)$ has at least one ideal vertex, and the convergence should only be from the complement of some domains of $(\mathcal{P}_H^c(l^k), \frac{k}{2}g_i)$ (corresponding to “degenerate” polygons which are close to segments) to the complement of some neighborhoods of the ideal vertices of $\mathcal{P}_E^c(\alpha)$.

Hyperbolic and de Sitter polygons. Similar natural metrics can be constructed on the space of convex hyperbolic polygons of given edge lengths, using Theorem B_H rather than Theorem B_S . However, the convergence to the spaces of Euclidean polygons should be considered for spaces of hyperbolic polygons with fixed angles, rather than fixed edge lengths. Fixing the angles of convex hyperbolic polygons is equivalent to fixing the edge lengths of the dual polygons in the de Sitter plane (see e.g. [Cox43, Cox57]).

For convex polygons in the de Sitter plane, Theorem B_{dS} (as seen in section 6) can also be used to define natural metrics on the space of de Sitter polygons with given edge lengths. Now, the sum of the lengths of the edges of a convex de Sitter polygon (which is dual to a hyperbolic polygon) is more than 2π ; the arguments used in the proof of Theorem 7.3 have analogs for the de Sitter plane, which seem to indicate that, as the sum of the edge lengths converges to 2π , the spaces of de Sitter polygons, with adequately normalized metrics, converge to a space of Euclidean polygons, with the metric defined by [BG92].

Acknowledgements

I would like to thank Philippe Eyssidieux, Vincent Guirardel, Sergiu Moroianu and Boris Springborn for some helpful discussions and interesting comments related to the content of this paper.

References

- [Ale58] A. D. Alexandrow. *Konvexe polyeder*. Akademie-Verlag, Berlin, 1958.
- [BG92] Christophe Bavard and Étienne Ghys. Polygones du plan et polyèdres hyperboliques. *Geom. Dedicata*, 43(2):207–224, 1992.
- [Cau13] A. L. Cauchy. Sur les polygones et polyèdres, second mémoire. *Journal de l'Ecole Polytechnique*, 19:87–98, 1813.
- [Cox43] H. S. M. Coxeter. A geometrical background for de Sitter’s world. *Amer. Math. Monthly*, 50:217–228, 1943.
- [Cox57] H. S. M. Coxeter. *Non-Euclidean geometry*. Mathematical Expositions, no. 2. University of Toronto Press, Toronto, Ont., 1957. 3rd ed.
- [Cox93] H. S. M. Coxeter. *The real projective plane*. Springer-Verlag, New York, third edition, 1993. With an appendix by George Beck, With 1 IBM-PC floppy disk (5.25 inch; DD).
- [Deh16] M. Dehn. Über den Starrheit konvexer Polyeder. *Math. Ann.*, 77:466–473, 1916.

- [DM86] P. Deligne and G. D. Mostow. Monodromy of hypergeometric functions and nonlattice integral monodromy. *Inst. Hautes Études Sci. Publ. Math.*, (63):5–89, 1986.
- [Euc02] Euclid. *Elements*. Green Lion Press, Santa Fe, NM, 2002. All thirteen books complete in one volume, The Thomas L. Heath translation, Edited by Dana Densmore.
- [Fil] F. Fillastre. Spaces of polygons to spaces of polyhedra following Bavard, Ghys and Thurston. *math.MG/0308187*.
- [Fil92] P. Filliman. Rigidity and the Alexandrov-Fenchel inequality. *Monatsh. Math.*, 113(1):1–22, 1992.
- [Isk00] I. Iskhakov. *On hyperbolic surface tessellations and equivariant spacelike convex polyhedral surfaces in Minkowski space*. PhD thesis, Ohio State University, 2000.
- [Kan90] Edgar Kann. Infinitesimal rigidity of almost-convex oriented polyhedra of arbitrary Euler characteristic. *Pacific J. Math.*, 144(1):71–103, 1990.
- [KM95] Michael Kapovich and John Millson. On the moduli space of polygons in the Euclidean plane. *J. Differential Geom.*, 42(2):430–464, 1995.
- [KM99] Michael Kapovich and John J. Millson. On the moduli space of a spherical polygonal linkage. *Canad. Math. Bull.*, 42(3):307–320, 1999.
- [LegII] A.-M. Legendre. *Eléments de géométrie*. Paris, 1793 (an II). Première édition, note XII, pp.321–334.
- [LS00] François Labourie and Jean-Marc Schlenker. Surfaces convexes fuchsiennes dans les espaces lorentziens à courbure constante. *Math. Annalen*, 316:465–483, 2000.
- [Mil71] A. D. Milka. The isoperimetry of polygons on the sphere. *Ukrain. Geometr. Sb.*, (10):49–50, 1971.
- [O’N83] B. O’Neill. *Semi-Riemannian Geometry*. Academic Press, 1983.
- [Pog56] A. V. Pogorelov. A new proof of rigidity of convex polyhedra. *Uspehi Mat. Nauk (N.S.)*, 11(5(71)):207–208, 1956.
- [RR00] Lucio Rodríguez and Harold Rosenberg. Rigidity of certain polyhedra in $\mathbf{R}^3$. *Comment. Math. Helv.*, 75(3):478–503, 2000.
- [Sch98] Jean-Marc Schlenker. Métriques sur les polyèdres hyperboliques convexes. *J. Differential Geom.*, 48(2):323–405, 1998.
- [Sch01] Jean-Marc Schlenker. Convex polyhedra in Lorentzian space-forms. *Asian J. of Math.*, 5:327–364, 2001.
- [Sie02] A. Siegel. A Dido problem as modernized by Fejes Tóth. *Discrete Comput. Geom.*, 27(2):227–238, 2002.
- [Ste42] J. Steiner. Sur le maximum et le minimum des figures dans le plan, sur la sphère, et dans l’espace en général. *J. Reine Angew. Math.*, 24:93–152, 190–250, 1842.
- [Sto68] J. J. Stoker. Geometrical problems concerning polyhedra in the large. *Comm. Pure Appl. Math.*, 21:119–168, 1968.
- [Thu98] William P. Thurston. Shapes of polyhedra and triangulations of the sphere. In *The Epstein birthday schrift*, volume 1 of *Geom. Topol. Monogr.*, pages 511–549 (electronic). Geom. Topol. Publ., Coventry, 1998.
- [Vol56] Yu. A. Volkov. On deformations of a convex polyhedral angle. *Uspehi Mat. Nauk (N.S.)*, 11(5(71)):209–210, 1956.